

## **Product datasheet for AM31881RP-N**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## Pdcd1 Hamster Monoclonal Antibody [Clone ID: J43]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: J43

Applications: FC

Recommended Dilution: Flow Cytometry.

Reactivity: Mouse
Host: Hamster
Isotype: IgG

Clonality: Monoclonal

**Immunogen:** PD-1 cDNA transfectant.

Donor: mouse ascites fluid.

Fusion Partner: Mouse myeloma P3U1.

**Specificity:** This Monoclonal Antibody reacts with Mouse PD-1 (Programmed Death-1).

**Formulation:** PBS containing 0.02% Sodium Azide as preservative and EIA grade BSA as a stabilizing protein

to bring total protein concentration to 4-5 mg/mL.

Label: PE

State: Liquid purified IgG fraction

Label: R - Phycoerythrin

**Concentration:** lot specific

**Purification:** Protein G Chromatography

Conjugation: PE

Storage: Store the antibody undiluted at 2-8°C.

**DO NOT FREEZE!** 

**Stability:** Shelf life: one year from despatch.

Gene Name: programmed cell death 1

Database Link: Entrez Gene 18566 Mouse

Q02242



## Pdcd1 Hamster Monoclonal Antibody [Clone ID: J43] - AM31881RP-N

Background:

CD279 is 50-55kD membrane protein which is a member of the CD28 family, and functions mainly as a negative regulator of T-cell activation. CD279 has two specific ligands; CD274 (PD-L1) and CD273 (PD-L2), and their interaction is key in the balance between stimulatory and inhibitory signals needed for effective immune responses to microbes and self-tolerance. CD279 is inducibly expressed by T-cells, B-cells, NK-T-cells and monocytes upon activation. Loss of CD279 function has been associated with a number of autoimmune diseases, including rheumatoid arthritis, type I diabetes and ankylosing spondylitis. Recent studies suggest that CD279 could be targeted therapeutically in the treatment of HIV infection to reduce T-cell exhaustion.

Synonyms: Protein PD-1, hPD-1, PDCD1

Note: TEST RESULTS (Figure 1.)

Tissue Distribution by Flow Cytometry Analysis:

Mouse Strain: Balb/c

Antibody Concentration Used: 2.0 µg/10e6 cells. Isotypic Control (shaded): Purified Hamster IgG.