

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for AM32091PU-N

## Fibronectin (FN1) (pFn) Mouse Monoclonal Antibody [Clone ID: BF12]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	BF12
Applications:	ELISA, IHC, WB
Recommended Dilution:	Western blotting: 1/1000. Immunohistochemistry on Frozen Sections. Other Immunoassays for cellular and tumor biology.
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Purified fibronectin from A8387 fibrosarcoma cells. Hybridoma produced by fusion between myeloma cells and Balb/c spleen cells.
Specificity:	The antibody reacts with the cellular and plasma forms of Fibronectin and recognizes the cell binding region.
Formulation:	PBS State: lg Fraction State: Liquid lg fraction Stabilizer: 1.0% BSA Preservative: 0.09% Sodium Azide
Concentration:	lot specific
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Gene Name:	fibronectin 1
Database Link:	<u>Entrez Gene 2335 Human</u> <u>P02751</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Fibronectin (FN1) (pFn) Mouse Monoclonal Antibody [Clone ID: BF12] – AM32091PU-N
Background:	Fibronectins are diverse high molecular weight cell adhesion glycoproteins found in basement membranes, in the interstitial connective tissue matrix, and in soluble in plasma and other body fluids. In adult tissues most fibronectin is produced by hepatocytes (pFn) that differ from locally produced cellular fibronectins. A monoclonal antibody (BF12) has been produced to fibronectin that reacts with both the cellular and plasma forms of the protein and recognizes the cell-binding region in fibronectin. The monoclonal antibody DH1 recognizes only cellular form of Fibronectin. DH1 is specific to the extra domain A (EDA) of the molecule. cFns are found in some basement membranes during development and only in distinct tissues in adult, for instance in smooth muscle and alveolar walls of airways. cFns are abundant in the plasma of carcinoma patients and in the stroma of various carcinomas.
Synonyms:	FN1, Cold-insoluble globulin, CIG

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US