

Product datasheet for AM32156PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Mucin-5AC Mouse Monoclonal Antibody [Clone ID: 9-13M1]

Product data:

Product Type: Primary Antibodies

Clone Name: 9-13M1

Applications: ELISA, IHC, R, WB

Recommended Dilution: Immunohistochemistry on cultured cells: Strongly positive on ethanol fixed epithelial cells.

Immunochemistry on Frozen Ethanol Fixed Sections.

Immunochemistry on Paraffin Sections: Very strong positive reaction with ethanol fixed

tissues.

Protease pretreatment is recommended for formalin post-fixed tissues.

Immunoblotting: Strongly positive without beta-mercaptoethanol pretreatment of Mucin

solution.

ELISA or IRMA in combination with the clone 9-13M1 or the mixture of anti-M1 Monoclonal

antibodies.

Reactivity: Human

Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: A BALB/c mouse was immunized with mucin isolated from an ovarian cyst fluid (pure

endocervical type according to the Fenoglio's classification). Splenocytes were fused with

mouse myeloma SP2/0 cells.

Specificity: This antibody clone *9-13M1* reacts with peptide core of Gastric Mucin (MUC5AC).

Formulation: PBS

State: Purified

State: Liquid purified IgG fraction Preservative: 0.05% Sodium Azide

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.





Mucin-5AC Mouse Monoclonal Antibody [Clone ID: 9-13M1] - AM32156PU-N

Gene Name: mucin 5AC, oligomeric mucus/gel-forming

Database Link: Entrez Gene 4586 Human

P98088

Background: Mucins are high molecular weight glycoproteins that are found especially in the secretions of

mucus membranes. Gastric Mucin 5AC antigen is found in columnar mucus cells of surface gastric epithelium and in goblet cells of the fetal and precancerous colon but not in normal colon. Resurgence of gastric mucin during colonic carcinogenesis is suggestive of either reexpression of the peptide core of gastric mucin in the adult colon or due to changes in the

glycosylation pattern of mucin, which expose the hidden Mucin 5AC antigen.

Synonyms: MUC5AC, MUC5, TBM, LeB