

## Product datasheet for **AM32233PU-N**

### TLR2 Mouse Monoclonal Antibody [Clone ID: mTL2.7]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	mTL2.7
Applications:	ELISA, IHC, IP
Recommended Dilution:	<b>Flow Cytometry.</b> The typical starting working dilution is 1/50. <b>Immunohistochemistry on frozen sections.</b> <b>Immunoassay.</b> <b>Immunoprecipitation.</b>
Reactivity:	Mouse
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Specificity:	Monoclonal antibody mT2.7 reacts with mouse Toll-like receptor 2 (TLR2). The monoclonal antibody mT2.7 stained overexpressed, as well as endogenous cell surface- and intercellular TLR2. The antibody does not affect cell activation through TLR2.
Formulation:	PBS State: Purified State: Liquid purified Ig fraction (0.2 µm filtered) Stabilizer: 0,1% BSA Preservative: 0.02% sodium azide
Concentration:	lot specific
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Gene Name:	toll like receptor 2
Database Link:	<a href="#">O60603</a>



[View online »](#)

**Background:**

Toll-like receptors (TLR) are highly conserved throughout evolution and have been implicated in the innate defense to many pathogens. In *Drosophila* toll is required for the anti-fungal response, while the related 18-wheeler is involved in antibacterial defenses. In mammals, TLR identified as type I transmembrane signaling receptors with pattern recognition capabilities, have been implicated in the innate host defense to pathogens. TLR2 has been identified as a receptor that is central to the innate immune response to lipoproteins of Gram-negative bacteria, several whole Gram-positive bacteria, as well as a receptor for peptidoglycan and lipoteichoic acid and other bacterial cell membrane products. A functional interaction between TLR2 and TLR6 in the cellular response to various bacterial products has been discovered. The currently accepted paradigm regards TLR2 as an essential receptor for many eubacterial cell wall components, including lipoproteins and peptidoglycan. Bacterial species as diverse as mycobacteria, spirochetes, mycoplasma, *Staphylococcus aureus*, and *Streptococcus pneumoniae* have all been shown to mediate cellular activation via TLR2.

**Synonyms:**

Toll-like receptor 2