

Product datasheet for **AM32313PU-N**

CD95 (FAS) Mouse Monoclonal Antibody [Clone ID: B-R18]

Product data:

Product Type:	Primary Antibodies
Clone Name:	B-R18
Recommended Dilution:	Flow Cytometry.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	BALB/c mice (Iffa Credo) were immunized with purified recombinant Fas antigen and fused with X63/Ag.8653 mouse myeloma cells.
Specificity:	This antibody clone <i>B-R18</i> specifically recognizes CD95. <i>B-R18</i> reacts with peripheral lymphocytes and especially with peripheral monocytes. It stains human B cell lines like pre-B cells, EBV cells, Burkitt cells and plasmacytoma cells. It also binds to human T cell lines, myeloid cell lines, hepatocyte carcinoma and endothelial cells.
Formulation:	PBS State: Purified State: Liquid purified IgG fraction Preservative: 0.05% Sodium Azide
Concentration:	lot specific
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Gene Name:	Fas cell surface death receptor
Database Link:	Entrez Gene 355 Human P25445



[View online »](#)

Background:

CD95, also known as FAS or APO1, is a 36 kDa cell surface type I membrane glycoprotein with an apparent molecular weight of 44 kDa on SDS PAGE. CD95 is a member of the TNF receptor family, which includes TNFR1, TNFR2, CD27, CD30 and CD40. Binding of CD95 Ligand to CD95 or crosslinking of CD95 by anti CD95 monoclonal antibodies leads to apoptosis of CD95 expressing cells. CD95 belongs to a subgroup of family members that have a death domain (DD) which contains 70 amino acids near the carboxyl terminal region of the molecule. The binding of adaptor molecules to this DD is responsible for transmitting the death signal for apoptosis. Stimulation of CD95 results in aggregation of its DD, leading to the recruitment of FADD and caspase 8 that together with the receptor form the death inducing signaling complex (DISC). CD95/CD95L is involved in the peripheral deletion of activated mature T cells at the end of the immune response and defects in this pathway predispose to autoimmune disorders. CD95 is also involved in killing of targets such as virus infected cells or cancer cells and killing of inflammatory cells at immune privileged sites.

Synonyms:

FASLG receptor, Apo-1 antigen, APT1, FAS1, TNFRSF6