

Product datasheet for **AM32426FC-N**

HLA Class II DR Mouse Monoclonal Antibody [Clone ID: B-C10]

Product data:

Product Type:	Primary Antibodies
Clone Name:	B-C10
Applications:	FC
Recommended Dilution:	Flow Cytometry: Use 10 µl to label 10 ⁶ cells or 100 µl of whole blood.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	RPMI 8866 and DAUDI cell lines. Myeloma X63/AG.8653 x Balb/c spleen cells.
Specificity:	This Monoclonal antibody recognises the HLA Class II (DR) membrane.
Formulation:	PBS Label: FITC State: Purified State: Liquid purified IgG fraction Stabilizer: 5% BSA Preservative: 0.09% Sodium Azide
Purification:	Ion Exchange Chromatography
Conjugation:	FITC
Storage:	Store undiluted at 2-8°C. DO NOT FREEZE!
Stability:	Shelf life: one year from despatch.




[View online »](#)

Background:

Major histocompatibility complex (MHC) class II molecules destined for presentation to CD4⁺ helper T cells is determined by two key events. These events include the dissociation of class II-associated invariant chain peptides (CLIP) from an antigen binding groove in MHC II alpha/beta dimers through the activity of MHC molecules HLA-DM and -DO, and subsequent peptide antigen binding. Accumulating in endosomal/lysosomal compartments and on the surface of B cells, HLA-DM, -DO molecules regulate the dissociation of CLIP and the subsequent binding of exogenous peptides to HLA class II molecules (HLA-DR, -DQ and -DP) by sustaining a conformation that favors peptide exchange. RFLP analysis of HLA-DM genes from rheumatoid arthritis (RA) patients suggests that certain polymorphisms are genetic factors for RA susceptibility. HLA-B belongs to the HLA class I heavy chain paralogs. Class I molecules play a central role in the immune system by presenting peptides derived from the endoplasmic reticulum lumen. HLA-B and -C can form heterodimers consisting of a membrane anchored heavy chain and a light chain (beta-2-Microglobulin). Polymorphisms yield hundreds of HLA-B and -C alleles.

Synonyms:

HLA-DR, HLA class II histocompatibility antigen DR, MHC class II antigen DR

Product images:

A typical staining pattern of lymphocytes with the B-C10 monoclonal antibody.