

Product datasheet for AM32966PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Psoriasin (S100A7) Mouse Monoclonal Antibody [Clone ID: 5C2]

Product data:

Product Type: Primary Antibodies

Clone Name: 5C2

Applications: ELISA, WB

Recommended Dilution: ELISA: This antibody is reactive to recombinant S100A7 protein.

Western blot: This antibody is reactive to 36.85KDa GST tagged recombinant S100A7 protein.

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: A short peptide (tdyhkqshgaapcsggs) derived from Human S100A7 (Psoriasin) conjugated with

KLH

Specificity: This antibody is reactive to S100A7 peptide (tdyhkqshgaapcsggs) and GST tagged

recombinant S100A7 protein.

Formulation: 0.01M PBS, pH 7.0 without preservatives

State: Purified

State: Lyophilized purified IgG fraction

Reconstitution Method: Restore with Double distillated water to adjust the final concentration to 1.0 mg/ml.

Concentration: 1.0 mg/ml (after reconstitution)

Purification: Affinity Chromatography on Protein G

Conjugation: Unconjugated

Storage: Prior to reconstitution store at 2-8°C.

Following reconstitution store undiluted at 2-8°C for one month

or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: S100 calcium binding protein A7

Database Link: Entrez Gene 6278 Human

P31151





Psoriasin (S100A7) Mouse Monoclonal Antibody [Clone ID: 5C2] - AM32966PU-N


Background: S100A7 is 11 kilo-dalton calcium binding protein with 2 EF motifs. It is also called psoriasin

due to its significantly elevated level in the skin lesion of psoriasis. The protein is over-expressed in squamous cell tumors of the skin, lung and breast. S100A7 expression is

induced after activation of epidermal growth factor receptor.

Synonyms: S100-A7, S100A7C, PSOR1

Product images:



The GST tagged recombinant S100A7 was loaded at 200 ng/lane. The anti-S100A7 clone 5C2 was used at 1 ug/ml, and the HRP conjugated Goat anti-Mouse IgG was used at 1/2,000 dilution.