

Product datasheet for AM33267PU-T

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD45 (PTPRC) (CD45RB) Mouse Monoclonal Antibody [Clone ID: SPM569]

Product data:

Product Type: Primary Antibodies

Clone Name: SPM569
Applications: FC, IF, IHC

Recommended Dilution: Flow Cytometry: $0.5-1 \mu g/106$ cells.

Immunofluorescence: 0.5-1 µg/ml.

Immunohistochemistry on Formalin-Fixed Paraffin Sections: 0.5-1 µg/ml for 30 minutes

at RT.

Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH

6.0, for 10-20 min followed by cooling at RT for 20 minutes. **Positive Control:** Ramos, U-698, or GA-10 cells, Tonsil.

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality:

Immunogen: Human peripheral blood lymphocytes maintained in T cell growth factor.

Specificity: This Monoclonal *SPM569* antibody recognizes Human CD4RB5. Other species not tested.

Cellular Localization: Cell surface and Cytoplasmic.

Formulation: 10mM PBS

State: Purified

Monoclonal

State: Liquid purified IgG fraction from Bioreactor Concentrate

Stabilizer: 0.05% BSA

Preservative: 0.05% Sodium Azide

Concentration: lot specific

Purification: Protein A/G Chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 180-220 kDa





Gene Name: protein tyrosine phosphatase, receptor type C

Database Link: Entrez Gene 5788 Human

P08575


Background: CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane

glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated

neoplasms.

Synonyms: PTPRC, Leukocyte common antigen, L-CA, T200

Product images:



Formalin-Fixed, Paraffin-Embedded Human Lymphoma stained with CD45RB Antibody (Clone SPM569).