

Product datasheet for **AM33286PU-T**

Hepatocyte Specific Antigen (HSA) Mouse Monoclonal Antibody [Clone ID: SPM582]

Product data:


Product Type:	Primary Antibodies
Clone Name:	SPM582
Applications:	IF, IHC
Recommended Dilution:	Immunofluorescence: 0.5-1 µg/ml. Immunohistochemistry on Formalin-Fixed Paraffin Sections: 1-2 µg/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 1mM EDTA, pH 7.5-8.5, for 10-20 min followed by cooling at RT for 20 minutes. Recommended Positive Control: Liver or Hepatocellular Carcinoma (HCC).
Reactivity:	Canine, Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Extract of a formalin-fixed, rejected-allograft of a Human liver.
Specificity:	Hepatocyte Specific Antigen, also called Hepatocyte Paraffin 1 or Hep Par 1, localizes to the mitochondria of hepatocytes. It is a sensitive marker for distinguishing hepatocellular carcinomas (HCC) from other metastatic carcinomas as well as cholangio-carcinomas. HCC's occur primarily in the stomach, but they are also found in many other organs. The Hepatocyte Specific Antigen may also be a useful marker for intestinal metaplasia. Reportedly, strong expression of the Hepatocyte Specific Antigen correlates with smaller tumor size and longer patient survival. Occasionally, Hepatocyte Specific Antigen is also found in gastric carcinomas as well as in a few other non-hepatic tumors. Cellular Localization: Finely granular cytoplasmic.
Formulation:	10mM PBS State: Purified State: Liquid purified IgG fraction from Bioreactor Concentrate Stabilizer: 0.05% BSA Preservative: 0.05% Sodium Azide
Concentration:	lot specific
Purification:	Protein A/G Chromatography



[View online »](#)

Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C. DO NOT FREEZE!
Stability:	Shelf life: one year from despatch.
Background:	Hepatoblastoma is the most common primary tumor of the liver in children. The use of specific hepatocyte markers and also of alpha Fetoprotein or carcinoembryonic antigen are useful for the identification of normal and malignant fetal hepatocytes.
Synonyms:	HepPar-1, hepatocellular carcinoma marker, hepatocyte marker

Product images:



Formalin-paraffin Human Hepatocellular Carcinoma stained with HSA Antibody Cat.-No AM33286PU (Clone SPM582).