

Product datasheet for AM33296PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Prolactin Receptor (PRLR) Mouse Monoclonal Antibody [Clone ID: B6.2+PRLR742]

Product data:

Product Type: Primary Antibodies

Clone Name: B6.2+PRLR742

Applications: FC, IF, IHC, IP, WB

Recommended Dilution: ELISA: Use BSA free Antibody for coating.

Flow Cytometry: 0.5-1 μg/million cells. **Immunofluorescence:** 0.5-1 μg/ml. **Western Blotting:** 0.25-0.5 μg/ml.

Immunoprecipitation: 0.5-1 μg/500 μg protein lysate.

Immunohistochemistry on Frozen and Formalin-Fixed Paraffin Sections: 0.5-1.0 µg/ml

for 30 minutes at RT.

Staining of formalin/paraffin tissues requires digestion of tissue sections with pepsin at

1mg/ml Tris-HCl, pH 2.0 for 15 min at RT or 10 min at 37°C.

Positive Control: Breast, placenta, liver or pancreas.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Semi-purified Human Prolactin Receptor (B6.2) and Recombinant Human Prolactin

Receptor (PRLR742).

Specificity: It recognizes a protein of 70kDa, identified as Prolactin Receptor.

Cellular Localization: Cell surface and cytoplasmic.

Formulation: 10mM PBS

State: Purified

State: Liquid purified IgG fraction from Bioreactor Concentrate

Stabilizer: 0.05% BSA

Preservative: 0.05% Sodium Azide

Concentration: lot specific

Purification: Protein A/G Chromatography

Conjugation: Unconjugated





Storage: Store undiluted at 2-8°C.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 70 kDa

Gene Name: prolactin receptor

Database Link: Entrez Gene 5618 Human

P16471

Background: Prolactin is a pituitary hormone involved in the stimulation of milk production, salt and water

regulation, growth, development and reproduction. The initial step in its action is the binding to a specific membrane receptor (prolactin receptor), which belongs to the superfamily of class 1 cytokine receptors. The function of the prolactin receptor is mediated, at least in part, by two families of signaling molecules: Janus kinases and signal transducers and activators of

transcription.

Synonyms: PRLR, PRL-R