

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AM33296PU-T

Prolactin Receptor (PRLR) Mouse Monoclonal Antibody [Clone ID: B6.2+PRLR742]

Product data:

Product Type:	Primary Antibodies
Clone Name:	B6.2+PRLR742
Applications:	FC, IF, IHC, IP, WB
Recommended Dilution:	 ELISA: Use BSA free Antibody for coating. Flow Cytometry: 0.5-1 µg/million cells. Immunofluorescence: 0.5-1 µg/ml. Western Blotting: 0.25-0.5 µg/ml. Immunoprecipitation: 0.5-1 µg/500 µg protein lysate. Immunohistochemistry on Frozen and Formalin-Fixed Paraffin Sections: 0.5-1.0 µg/ml for 30 minutes at RT. Staining of formalin/paraffin tissues requires digestion of tissue sections with pepsin at 1mg/ml Tris-HCl, pH 2.0 for 15 min at RT or 10 min at 37°C. Positive Control: Breast, placenta, liver or pancreas.
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Semi-purified Human Prolactin Receptor (B6.2) and Recombinant Human Prolactin Receptor (PRLR742).
Specificity:	lt recognizes a protein of 70kDa, identified as Prolactin Receptor. Cellular Localization : Cell surface and cytoplasmic.
Formulation:	10mM PBS State: Purified State: Liquid purified IgG fraction from Bioreactor Concentrate Stabilizer: 0.05% BSA Preservative: 0.05% Sodium Azide
Concentration:	lot specific
Purification:	Protein A/G Chromatography
Conjugation:	Unconjugated



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

ORIGENE Prolactin Receptor (PRLR) Mouse Monoclonal Antibody [Clone ID: B6.2+PRLR742] – AM33296PU-T	
Storage:	Store undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	70 kDa
Gene Name:	prolactin receptor
Database Link:	<u>Entrez Gene 5618 Human</u> <u>P16471</u>
Background:	Prolactin is a pituitary hormone involved in the stimulation of milk production, salt and water regulation, growth, development and reproduction. The initial step in its action is the binding to a specific membrane receptor (prolactin receptor), which belongs to the superfamily of class 1 cytokine receptors. The function of the prolactin receptor is mediated, at least in part, by two families of signaling molecules: Janus kinases and signal transducers and activators of transcription.
Synonyms:	PRLR, PRL-R

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US