

Product datasheet for AM33360PU-T

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD46 Mouse Monoclonal Antibody [Clone ID: 122.2]

Product data:

Product Type: Primary Antibodies

Clone Name: 122.2

Applications: FC, FN, IF, IHC, IP, WB

Recommended Dilution: Functional Assays (Neutralization): Use Azide free Antibody.

Western Blot: 0.5-1 µg/ml.

Flow Cytometry: $0.5-1 \mu g/106$ cells.

Immunoprecipitation: 1-2 μg/500 μg protein lysate.

Immunofluorescence: 1-2 µg/ml.

Immunohistochemistry on Frozen Sections: 0.5-1 µg/ml for 30 minutes at RT.

Recommended Positive Control: HeLa, K-562 or MOLT-4 cells, Kidney.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Recombinant Human CD46 protein.

Specificity: This antibody recognizes Human CD46. Other Species not tested.

Cellular Localization: Cell surface.

Formulation: 10mM PBS

State: Purified

State: Liquid purified IgG fraction from Bioreactor Concentrate

Stabilizer: 0.05% BSA

Preservative: 0.05% Sodium Azide

Concentration: lot specific

Purification: Protein A/G Chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 56-66 kDa





Gene Name: CD46 molecule

Database Link: Entrez Gene 4179 Human

P15529


Background: CD46 acts as a cofactor for complement factor I, a serine protease which protects autologous

cells against complement-mediated injury by cleaving C3b and C4b deposited on host tissue. It may be involved in the fusion of the spermatozoa with the oocyte during fertilization. CD46 acts as a co-stimulatory factor for T-cells which induces the differentiation of CD4+ into T-regulatory 1 cells. T-regulatory 1 cells suppress immune responses by secreting interleukin-10, and therefore are thought to prevent autoimmunity. A number of viral and bacterial pathogens seem to exploit this property and directly induce an immunosuppressive

phenotype in T-cells by binding to CD46.

Synonyms: TLX, MIC10

Product images:



FCM staining of human PBMCs using CD46 Antibody (Clone 122.2).