

Product datasheet for **AM33366PU-T**

CDC20 Mouse Monoclonal Antibody [Clone ID: AR12]

Product data:

| | |
|-----------------------|--|
| Product Type: | Primary Antibodies |
| Clone Name: | AR12 |
| Applications: | FC, IF, IHC, IP, WB |
| Recommended Dilution: | ELISA: Use BSA free Antibody for coating. Flow Cytometry: 0.5-1 µg/million cells. Immunofluorescence: 0.5-1 µg/ml. Western Blotting: 0.5-1 µg/ml. Immunoprecipitation: 0.5-1 µg/500 µg protein lysate. Immunohistochemistry on Frozen and Formalin-Fixed Paraffin Sections: 0.5-1 µg/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes. Positive Control: Ramos or HeLa cells; Tonsil. |
| Reactivity: | Human |
| Host: | Mouse |
| Isotype: | IgG1 |
| Clonality: | Monoclonal |
| Immunogen: | Urea-denatured His6 Cdc20 Human recombinant protein. |
| Specificity: | This antibody reacts with Human Cdc20 (55 kDa). Other species not tested. Cellular Localization: Cytoplasmic. |
| Formulation: | 10mM PBS State: Purified State: Liquid purified IgG fraction from Bioreactor Concentrate Stabilizer: 0.05% BSA Preservative: 0.05% Sodium Azide |
| Concentration: | lot specific |
| Purification: | Protein A/G Chromatography |
| Conjugation: | Unconjugated |
| Storage: | Store undiluted at 2-8°C. |



[View online »](#)

| | |
|--------------------------------|--|
| Stability: | Shelf life: one year from despatch. |
| Predicted Protein Size: | 55 kDa |
| Gene Name: | cell division cycle 20 |
| Database Link: | Entrez Gene 991 Human Q12834 |
| Background: | Cyclins, regulatory subunits which associate with kinases, control many of the important steps in cell cycle progression. The Cdc2 protein kinase (p34Cdc2) exhibits protein kinase activity <i>in vitro</i> and exists in a complex with both cyclin B and a protein homologous to p13SUC1. Cdc2 kinase is the active subunit of the M phase promoting factor (MPF) and the M phase-specific Histone H1 kinase. The p34Cdc2/cyclin B complex is required for the G2 to M transition. An additional cell cycle-dependent protein kinase, termed p55cdc, exhibits a high degree of homology with the <i>S. cerevisiae</i> proteins Cdc20 and Cdc4. The p55cdc transcript is readily detectable in a variety of cultured cell lines in growth phase, but disappears when cell growth is chemically arrested. |
| Synonyms: | p55CDC |