

Product datasheet for AM33412PU-N

PD1 (PDCD1) Mouse Monoclonal Antibody [Clone ID: 7C9]

Product data:

Product Type: Primary Antibodies Clone Name: 709 **Applications:** ELISA, WB Recommended Dilution: Indirect ELISA. Western Blot. **Reactivity:** Human Mouse Host: Isotype: lgG1 **Clonality:** Monoclonal Immunogen: 40kDa recombinant Human Programmed Cell Death-1 (PD-1). Specificity: Reactive with recombinant Human Programmed Cell Death-1 (PD-1). Reactivity with natural PD-1 has not been evaluated. Formulation: 0.01M PBS, pH 7.2 without preservatives State: Aff - Purified State: Lyophilized purified IgG fraction **Reconstitution Method:** Restore with Double distillated water to adjust the final concentration to 1.0 mg/ml. **Purification:** Affinity Chromatography on Protein G **Conjugation:** Unconjugated Storage: Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing. Stability: Shelf life: one year from despatch. Gene Name: programmed cell death 1 Database Link: Entrez Gene 5133 Human <u>Q15116</u>



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	PD1 (PDCD1) Mouse Monoclonal Antibody [Clone ID: 7C9] – AM33412PU-N
Background:	Programmed Cell Death-1 (PD-1), also called CD279, is a 288 amino acid cell surface protein that belongs to the immunoglobulin superfamily. The protein is expressed on T cells, B cells and macrophages. When PD-1 is joined with its ligand (CD274 or B7-H1), the immune cell's activation and effector function are attenuated. Research suggested that PD1 pathway is mainly involved with attenuation at periphery sites while the B7/CTLA-4 pathway is primarily involved with the attenuation of the activation of immune cells in secondary lymphoid organs. PD-1 pathway is implicated with cancer's immune-evasion, autoimmunity, and <i>Listeria</i> <i>monocytogenes</i> infection.

Synonyms: Protein PD-1, hPD-1, PDCD1

Product images:



Western blot analysis showing 1 ug/well of recombinant Human PD-1 detected with 1 ug/ml of CD279 / PD1 Antibody (Clone 7C9).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US