

Product datasheet for AM50012PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PIG3 (TP53I3) (1-332) Mouse Monoclonal Antibody [Clone ID: AT1C9]

Product data:

Product Type: Primary Antibodies

Clone Name: AT1C9

Applications: ELISA, FC, WB

Recommended Dilution: ELISA.

Western blot (1:3,000).

Flow cytometry.

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Recombinant human TP53I3 (1-332aa) purified from *E. coli*

Specificity: This antibody detects Recombinant human TP53I3 (1-332aa).

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein-G affinity chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: tumor protein p53 inducible protein 3

Database Link: Entrez Gene 9540 Human

Q53FA7



PIG3 (TP53I3) (1-332) Mouse Monoclonal Antibody [Clone ID: AT1C9] – AM50012PU-N


Background:

TP53I3 participates in the generation of reactive oxygen species (ROS). TP53I3 has low NADPH-dependent naphtoquinone reductase activity, with a preference for 1,2-naphtoquinone over 1,4-naphtoquinone. TP53I3 has low NADPH-dependent diamine reductase activity (in vitro). TP53I3 is localized to the cytoplasm and induced in primary, non-transformed and transformed cell cultures after exposure to genotoxic agents. TP53I3 microsatellite polymorphism is associated with differential susceptibility to cancer.

Synonyms:

p53-induced gene 3 protein, PIG3

Product images:



The extracts of PC-3 (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human KLK3 antibody (1:500 ~ 1:5000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.