

Product datasheet for AM50026PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

PFKM Mouse Monoclonal Antibody [Clone ID: AT2F11]

Product data:

Product Type: Primary Antibodies

Clone Name: AT2F11

Applications: ELISA, FC, WB

Recommended Dilution: ELISA.

Western blot: 1/500-1/2000.

Recommended Sarting Dilution: 1/2000.

Flow Cytometry.

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Recombinant Human PFKM (1-780aa) purified from *E. coli*

Specificity: This Monoclonal antibody recognizes Human PFKM. Other species not tested.

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Affinity Chromatography on Protein A

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: phosphofructokinase, muscle

Database Link: <u>Entrez Gene 5213 Human</u>

P08237



PFKM Mouse Monoclonal Antibody [Clone ID: AT2F11] - AM50026PU-N

Background: PFKM (phosphofructokinase, muscle) is a regulatory glycolytic enzyme that convert fructose

6-phosphate and ATP into fructose 1,6-bisphosphate (through PFK-1), fructose 2,6-bisphosphate (through PFK-2) and ADP. Three phosphofructokinase isozymes exist in humans: muscle, liver and platelet. Mutations in this gene have been associated with

glycogen storage disease type VII, also known as Tarui disease.

Synonyms: Phosphofructokinase-M, Phosphohexokinase, Phosphofructokinase 1, PFK1, PFK-1, PFK-A,


PFKX, GSD7

Protein Families: Druggable Genome

Protein Pathways: Fructose and mannose metabolism, Galactose metabolism, Glycolysis / Gluconeogenesis,

Metabolic pathways, Pentose phosphate pathway

Product images:



The HepG2 cell lysate (30ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-Human PFKM antibody (1/500~1/2000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.