

Product datasheet for AM50044PU-S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

HEXA (89-529) Mouse Monoclonal Antibody [Clone ID: AT20F1]

Product data:

Product Type: Primary Antibodies

Clone Name: AT20F1

Applications: ELISA, FC, WB

Recommended Dilution: ELISA.

Western blot: Recommended starting dilution is 1:3000.

Flow cytometry.

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Recombinant human HEXA (89-529aa) purified from *E. coli*

Specificity: This antibody detects HEXA at aa 89-529.

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein-A affinity chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: hexosaminidase subunit alpha

Database Link: Entrez Gene 3073 Human

P06865






Background:

HEXA (Hexosaminidase A), also designated beta-Hexosaminidase A, is responsible for the degradation of GM2 gangliosides, and a variety of other molecules containing terminal N-acetyl hexosamines, in the brain and other tissues. A mutation in the a subunit of hexosaminidase is the cause of Tay-Sachs disease (TSD), also known as GM2-gangliosidosis type I. TSD is a fatal autosomal recessive lysosomal storage disease of the central nervous system (CNS) caused by insufficient activity of the HEXA enzyme that results in a failure to process GM2 gangliosides. The accumulation of GM2 ganglioside in the absence of HEXA activity causes progressive destruction of the CNS.

Synonyms:

beta-N-acetylhexosaminidase; MGC99608; N-acetyl-beta-glucosaminidase; TSD

Product images:



The cell lysates of MCF7 (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human HEXA antibody (1:3000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.