

Product datasheet for AM50046PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CIAO1 Mouse Monoclonal Antibody [Clone ID: AT6C9]

Product data:

Product Type: Primary Antibodies

Clone Name: AT6C9

Applications: ELISA, WB

Recommended Dilution: The antibody has been tested by ELISA, Western blot analysis to assure specificity and

reactivity. Since application varies, however, each investigation should be titrated by the reagent to obtain optimal results. Recommended dilution range for Western blot analysis is

1:500 ~ 1:5000. Recommended starting dilution is 1:1000.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Recombinant human CIAO1 (1-339aa) purified from E. coli

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein-A affinity chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: cytosolic iron-sulfur assembly component 1

Database Link: Entrez Gene 9391 Human

076071





Background: CIAO1 is a essential component of the cytosolic iron-sulfur (Fe/S) protein assembly

machinery. It is a multiprotein complex that mediates the incorporation of iron-sulfur cluster

into extramitochondrial Fe/S proteins. CIAO1 seems to specifically modulate the


transactivation activity of WT1 and may function to regulate the physiological functions of WT1 in cell growth and differentiation. As part of the mitotic spindle-associated MMXD

complex, it may play a role in chromosome segregation.

Synonyms: CIA1; WDR39

Protein Families: Transcription Factors

Product images:



The cell lysate of Jurkat (30ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CIAO1 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.