

Product datasheet for AM50075PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ING2 Mouse Monoclonal Antibody [Clone ID: AT39E5]

Product data:

Product Type: Primary Antibodies

Clone Name: AT39E5

Applications: ELISA, WB

Recommended Dilution: The antibody has been tested by ELISA, Western blot analysis to assure specificity and

reactivity. Since application varies, however, each investigation should be titrated by the

reagent to obtain optimal results. Recommended starting dilution is 1:1000.

Reactivity: Human Host: Mouse

Clonality: Monoclonal

Immunogen: Recombinant human ING2 (1-280aa) purified from E. coli

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein-A affinity chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: inhibitor of growth family member 2

Database Link: Entrez Gene 3622 Human

Q9H160

ING2 Mouse Monoclonal Antibody [Clone ID: AT39E5] - AM50075PU-N

Background: ING2 is a member of the inhibitor of growth (ING) family. Members of the ING family


associate with and modulate the activity of histone acetyltransferase (HAT) and histone deacetylase (HDAC) complexes and function in DNA repair and apoptosis. This protein seems to be involved in p53/TP53 activation and p53/TP53-dependent apoptotic pathways, probably by enhancing acetylation of p53/TP53. Also, It is component of a mSin3A-like corepressor complex, which is probably involved in deacetylation of nucleosomal histones. ING2 activity

seems to be modulated by binding to phosphoinositides.

Synonyms: p33ING2, Inhibitor of growth protein 2, Inhibitor of growth 1-like protein

Protein Families: Druggable Genome, Transcription Factors

Product images:



The cell lysates of HeLa (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human ING2 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.