

Product datasheet for AM50081PU-S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

CKMT1B Mouse Monoclonal Antibody [Clone ID: AT17A2]

Product data:

Product Type: Primary Antibodies

Clone Name: AT17A2

Applications: ELISA, FC, IF, WB

Recommended Dilution: ELISA.

Western blot. Flow Cytometry.

Immunofluorescence.

Recommended starting dilution is 1:1000.

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Recombinant human CKMT1A (40-417aa) purified from E. coli

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein-A affinity chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: creatine kinase, mitochondrial 1B

Database Link: Entrez Gene 548596 HumanEntrez Gene 1159 Human

P12532



CKMT1B Mouse Monoclonal Antibody [Clone ID: AT17A2] - AM50081PU-S

Background: CKMT1A, also known as Creatine kinase, mitochondrial 1A, is 417 amino acid protein.

Mitochondrial creatine kinase (MtCK) exists as two isoenzymes, sarcomeric MtCK (CKMT2) and ubiquitous MtCK (CKMT1). It is responsible for the transfer of high energy phosphate from mitochondria to the cytosolic carrier, creatine. Many malignant cancers with poor prognosis have shown overexpression of ubiquitous mitochondrial creatine kinase; this may


be related to high energy turnover and failure to eliminate cancer cells via apoptosis.

Synonyms: CKMT1A, CKMT1B, U-MtCK, Mia-CK, Creatine kinase U

Protein Families: Druggable Genome

Protein Pathways: Arginine and proline metabolism, Metabolic pathways

Product images:



The cell lysates of HeLa (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CKMT1A antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.