

Product datasheet for AM50130PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cytokeratin 14 (KRT14) Mouse Monoclonal Antibody [Clone ID: SPM263]

Product data:

Product Type: Primary Antibodies

Clone Name: SPM263

Applications: FC, IF, IHC, IP, WB

Recommended Dilution: ELISA: Use BSA free Antibody for coating.

Flow Cytometry: 0.5-1 μg/million cells. **Immunofluorescence:** 1-2 μg/ml. **Western Blotting:** 0.25-0.5 μg/ml.

Immunoprecipitation: 1-2 μg/500 μg protein lysate.

Immunohistochemistry on Frozen and Formalin-fixed Sections: 0.5-1.0 µg/ml for

30 minutes at RT.

Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, pH

6.0, for 10-20 min followed by cooling at RT for 20 minutes. *Positive Control*: A431 cells, Skin or Squamous cell carcinoma.

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG3

Clonality: Monoclonal

Immunogen: A synthetic peptide of 15 amino acid from the C-terminus of human keratin 14.

Specificity: Cytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight

keratins and exists in combination with keratin 5 (type II or B or basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing between basal and non-basal subtypes of breast carcinomas. Anti-CK14 is also a good marker for differentiation of intraductal from invasive salivary duct carcinoma by the positive staining of basal cells surrounding the in-situ neoplasm as well as for differentiation of benign prostate from prostate carcinoma. Furthermore, this antibody has been useful in separating oncocytic tumors of the kidney from its renal mimics, and in identifying metaplastic carcinomas of the

breast.

Cellular Localization: Cytoplasmic.





Formulation: 10mM PBS

State: Purified

State: Liquid purified IgG fraction from Bioreactor Concentrate

Stabilizer: 0.05% BSA

Preservative: 0.05% Sodium Azide

Concentration: lot specific

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 50 kDa

Gene Name: keratin 14

Database Link: Entrez Gene 3861 Human

P02533

Synonyms: Cytokeratin-14, CK14, Keratin 14, K14, KRT14

Product images:



Formalin-Paraffin prostate (10X) stained with Cytokeratin 14 Antibody (Clone SPM263).





Formalin-Paraffin prostate (20X) stained with Cytokeratin 14 Antibody (Clone SPM263).