

Product datasheet for AM50144PU-T

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MYOD1 Mouse Monoclonal Antibody [Clone ID: SPM427]

Product data:

Product Type: Primary Antibodies

Clone Name: SPM427

Applications: FC, IF, IHC, IP, WB

Recommended Dilution: ELISA: Use BSA free Antibody for coating.

Flow Cytometry: 0.5-1 μg/million cells. **Immunofluorescence:** 0.5-1 μg/ml. **Western Blotting:** 0.25-0.5 μg/ml.

Immunoprecipitation: 0.5-1 μg/500 μg protein lysate.

Immunohistochemistry on Frozen and Formalin-Fixed Paraffin Sections: 0.5- $1.0~\mu g/ml$

for 30 minutes at RT.

Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 1mM EDTA Buffer,

pH 7.5-8.5, for 10-20 min followed by cooling at RT for 20 minutes.

Positive Control: Rhabdomyosarcoma.

Reactivity: Chicken, Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Recombinant mouse MyoD1 protein.

Specificity: Recognizes a phosphor-protein of 45kDa, identified as MyoD1. The epitope of this

Monoclonal Antibody maps between amino acid 180-189 in the C-terminal of Mouse MyoD1

protein. It does not cross react with Myogenin, Myf5, or Myf6.

Antibody to MyoD1 labels the nuclei of myoblasts in developing muscle tissues. MyoD1 is not

detected in normal adult tissue, but is highly expressed in the tumor cell nuclei of

rhabdomyosarcomas. Occasionally nuclear expression of MyoD1 is seen in

ectomesenchymoma and a subset of Wilm's tumors. Weak cytoplasmic staining is observed

in several non-muscle tissues, including glandular epithelium and also in

rhabdomyosarcomas, neuroblastomas, Ewing's sarcomas and alveolar soft part sarcomas. *Cellular Localization*: Nuclear. *Only nuclear staining should be considered as evidence of*

skeletal muscle differentiation.





MYOD1 Mouse Monoclonal Antibody [Clone ID: SPM427] - AM50144PU-T

Formulation: 10mM PBS

State: Purified

State: Liquid purified IgG fraction from Bioreactor Concentrate

Stabilizer: 0.05% BSA

Preservative: 0.05% Sodium Azide

Concentration: lot specific

Purification: Protein A/G Chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 45 kDa


Gene Name: myogenic differentiation 1

Database Link: Entrez Gene 4654 Human

P15172

Synonyms: MYF3, MYOD, Myoblast determination protein 1, Myogenic factor 3, BHLHC1

Product images:



Formalin-paraffin rhabdomyosarcoma stained with MyoD1 Antibody (Clone SPM427).