

## Product datasheet for AM50175PU-T

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### TTF1 (NKX2-1) Mouse Monoclonal Antibody [Clone ID: SPM150]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: SPM150

**Applications:** FC, IF, IHC, IP, WB

**Recommended Dilution: ELISA:** Use BSA free Antibody for coating.

**Flow Cytometry:** 0.5-1 μg/million cells. **Immunofluorescence:** 0.5-1 μg/ml. **Western Blotting:** 0.5-1 μg/ml.

**Immunoprecipitation:** 1-2 μg/500 μg protein lysate.

Immunohistochemistry on Frozen and Formalin-Fixed Sections: 0.5-1 µg/ml for 30

minutes at RT.

Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH

6.0, for 10-20 min followed by cooling at RT for 20 minutes.

Positive Control: MAD109, MLE-15, H441-4, or H345 cells. Normal thyroid or lung.

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Rat full length TTF-1 recombinant protein.

**Specificity:** Recognizes a protein of 40kDa, identified as Thyroid transcription factor-1 (TTF-1).

TTF-1 is a member of the NKx2 family of homeodomain transcription factors. It is expressed in epithelial cells of the thyroid gland and the lung. Nuclei from liver, stomach, pancreas, small intestine, colon, kidney, breast, skin, testes, pituitary, prostate, and adrenal glands are unreactive. Anti-TTF-1 is useful in differentiating primary adenocarcinoma of the lung from metastatic carcinomas originating in the breast, mediastinal germ cell tumors, and malignant mesothelioma. It can also be used to differentiate small cell lung carcinoma from lymphoid infiltrates. Loss of TTF-1 expression in non-small cell lung carcinoma has been associated with aggressive behavior of such neoplasms. TTF-1 reactivity is also seen in thyroid malignancies.

Cellular Localization: Nuclear.



#### TTF1 (NKX2-1) Mouse Monoclonal Antibody [Clone ID: SPM150] - AM50175PU-T

Formulation: 10mM PBS

State: Purified

State: Liquid purified IgG fraction from Bioreactor Concentrate

Stabilizer: 0.05% BSA

Preservative: 0.05% Sodium Azide

**Concentration:** lot specific

**Purification:** Protein A/G Chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 40 kDa


Gene Name: NK2 homeobox 1

**Database Link:** Entrez Gene 7080 Human

P43699


Synonyms: NKX2-1, Nkx-2.1, TITF1, Thyroid nuclear factor 1, Thyroid transcription factor 1, TTF-1

# **Product images:**



Formalin-Paraffin lung adenocarcinoma (10X) stained with TTF-1 Antibody (Clone SPM150).





Formalin-Paraffin lung adenocarcinoma (20X) stained with TTF-1 Antibody (Clone SPM150).