

Product datasheet for AM50180PU-T

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Thymidylate Synthase (TYMS) Mouse Monoclonal Antibody [Clone ID: SPM453]

Product data:

Product Type: Primary Antibodies

Clone Name: SPM453

Applications: FC, IF, IHC, IP, WB

Recommended Dilution: ELISA: Use BSA free Antibody for coating.

Flow Cytometry: 0.5-1 μg/million cells. **Immunofluorescence:** 0.5-1 μg/ml. **Western Blotting:** 0.25-0.5 μg/ml.

Immunoprecipitation: 0.5-1 μg/500 μg protein lysate.

Immunohistochemistry on Frozen and Formalin-Fixed Sections: $0.5-1 \mu g/ml$ for 30 minutes at RT. Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.

Positive Control: 5-FU-resistant colon carcinoma cell lines (NCI H630R10, NCI H630R1); 5-FU-resistant breast cancer cell lines, MCF-Ad5 and MCF-Ad10. Colorectal, gastric, head & neck,

and breast carcinomas.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Recombinant human thymidylate synthase.

Specificity: It recognizes a protein of 36kDa, identified as Thymidylate Synthase (TS) (EC 2.1.1.45).

Cellular Localization: Cytoplasmic.

Formulation: 10mM PBS

State: Purified

State: Liquid purified IgG fraction from Bioreactor Concentrate

Stabilizer: 0.05% BSA

Preservative: 0.05% Sodium Azide

Concentration: lot specific

Purification: Protein A/G Chromatography

Conjugation: Unconjugated





Thymidylate Synthase (TYMS) Mouse Monoclonal Antibody [Clone ID: SPM453] - AM50180PU-T

Storage: Store undiluted at 2-8°C.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 36 kDa

Gene Name: thymidylate synthetase

Database Link: Entrez Gene 7298 Human

P04818

Background: TS converts deoxyuridine monophosphate (dUMP) to deoxythymidine monophosphate

(dTMP), which is essential for DNA biosynthesis. TS is also a critical target for the

fluoropyrimidines, an important group of antineoplastic drugs that are widely used in the treatment of solid tumors. Both 5-FU and fluorodeoxyuridine are converted in tumor cells to FdUMP which inactivates TS by formation of a ternary covalent complex in the presence of the folate cofactor 5,10-methylenetetrahydrofolate. Expression of TS protein is associated with response to 5-fluorouracil (5-FU) in human colorectal, gastric, head and neck, and breast

carcinomas.

Synonyms: TYMS, TSase, OK/SW-cl.29