

Product datasheet for **AM50186PU-S**

Vimentin (VIM) Mouse Monoclonal Antibody [Clone ID: SPM576]


Product data:

Product Type:	Primary Antibodies
Clone Name:	SPM576
Applications:	FC, IF, IHC, IP, WB
Recommended Dilution:	<p>ELISA: Use BSA free Antibody for coating.</p> <p>Flow Cytometry: 0.5-1 µg/million cells</p> <p>Immunofluorescence: 1-2 µg/ml.</p> <p>Western Blotting: 0.5-1 µg/ml.</p> <p>Immunoprecipitation: 1-2 µg/500 µg protein lysate.</p> <p>Immunohistochemistry on Frozen and Formalin-Fixed Paraffin Sections: 0.5-1 µg/ml for 30 minutes at RT.</p> <p>Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>Positive Control: Jurkat cells, Sarcomas, Melanomas.</p>
Reactivity:	Bovine, Canine, Chicken, Feline, Goat, Human, Porcine
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human vimentin recombinant protein.
Specificity:	<p>This MAb reacts with a 58kDa protein identified as vimentin. It shows no cross-reaction with other closely related intermediate filament proteins (IFP's) such as Desmin, Keratin, neurofilament, and glial fibrillary acid protein. Anti-Vimentin alone is of limited value as a diagnostic tool; however, when used in panels with other antibodies, it is useful for the sub-classification of a given tumor. Expression of vimentin, when used in conjunction with anti-keratin, is helpful when distinguishing melanomas from undifferentiated carcinomas and large cell lymphomas. All melanomas and Schwannomas react strongly with anti-vimentin. It labels a variety of mesenchymal cells, including melanocytes, lymphocytes, endothelial cells, and fibroblasts. Non-reactivity of anti-vimentin is often considered more useful than its positive reactivity, since there are a few tumors that do not contain vimentin, e.g. hepatoma and seminoma. Anti-vimentin is also useful as a tissue process control reagent.</p> <p>Cellular Localization: Cytoplasmic.</p> <p>Negative Species: Mouse and Rat.</p>



[View online »](#)

Formulation:	10mM PBS State: Purified State: Liquid purified IgG fraction from Bioreactor Concentrate Stabilizer: 0.05% BSA Preservative: 0.05% Sodium Azide
Concentration:	lot specific
Purification:	Protein A/G Chromatography
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	57-60 kDa
Gene Name:	vimentin
Database Link:	Entrez Gene 7431 Human P08670
Synonyms:	VIM

Product images:


Formalin-paraffin human melanoma stained with Vimentin Antibody (Clone SPM576).