

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AM50264PU-T

TMEM16A (ANO1) Mouse Monoclonal Antibody [Clone ID: DG1/447 + DOG-1.1]

Product data: Product Type: Primary Antibodies Clone Name: DG1/447 + DOG-1.1 **Applications:** IHC **Recommended Dilution: Immunohistochemistry on Formalin-fixed Sections:** 1.0 - 2.0 µg/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes. Positive Control: Gastrointestinal Stromal Tumor (GIST) or testicular germ cell tumor. Melanocytes in the basal layer of the epidermis and mast cells in the dermis of normal skin. **Reactivity:** Human Host: Mouse lgG1 Isotype: **Clonality:** Monoclonal Immunogen: Recombinant Human DOG-1 protein (DG1/447) and a synthetic peptide from Human DOG-1 protein (MSDFVDWVIPDIPKDISQQIHK-EKVLMVELFMREEQDKQQLLETCMEKER QKDEPPCNHHNTKACPDSLGSP-APSHAYHGGVL), conjugated to a carrier protein (DOG-1.1). Specificity: Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GIST's), c-kit signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be difficult to diagnose. Immunoreactivity for DOG-1 has been reported in 97.8 percent of scorable GIST's, including all c-kit negative GIST's. Overexpression of DOG-1 has been suggested to aid in the identification of GISTs, including Platelet-Derived Growth Factor Receptor Alpha mutants that fail to express c-kit antigen. The overall sensitivity of DOG1 and c-kit in GIST's is nearly identical: 94.4% vs. 94.7%. *Cellular Localization*: Cell Surface and Cytoplasmic. Formulation: 10mM PBS State: Purified State: Liquid purified IgG fraction from Bioreactor Concentrate Stabilizer: 0.05% BSA Preservative: 0.05% Sodium Azide

Purification:

lot specific

Protein A/G Chromatography

Concentration:




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	16A (ANO1) Mouse Monoclonal Antibody [Clone ID: DG1/447 + DOG-1.1] – AM50264PU-T
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~114 kDa
Gene Name:	anoctamin 1
Database Link:	<u>Entrez Gene 55107 Human</u> <u>Q5XXA6</u>
Background:	DOG1 gene, a gastrointestinal stromal tumor (GIST) specific gene, encoding for the hypothetical protein FLJ10261, which was named <i>Discovered on GIST 1</i> (DOG1). DOG1 protein is expressed ubiquitously in gastrointestinal stromal tumors irrespective of KIT or PDGFR alpha mutation status.
_	

Synonyms:

Anoctamin-1, Transmembrane protein 16A, DOG-1, ORAOV2, TAOS2, TMEM16A

Product images:



Formalin-fixed, paraffin-embedded human GIST (20X) stained with DOG1 Antibody AM50156PU (Clone DG1/447 + DOG1.1).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US