

Product datasheet for **AM50266PU-T**

EGFR Mouse Monoclonal Antibody [Clone ID: H9B4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	H9B4
Applications:	IP, WB
Recommended Dilution:	ELISA: Use BSA free Antibody for coating. Immunoprecipitation: 1-2 µg/500ug protein lysate. Western Blotting: 0.5-1.0 µg/ml for 2 hours at RT. Positive Control: A431 cells, Breast or Bladder cancer.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Purified EGFR from A431 cells.
Specificity:	This MAb reacts with a cytoplasmic domain of EGFR. EGFR is a type I receptor tyrosine kinase with sequence homology to erbB-1, -2, -3 -4 or HER-1, -2, -3 -4. It binds to Epidermal Growth Factor (EGF), Transforming Growth Factor-α (TGF-α), Heparin-binding EGF (HB-EGF), amphiregulin, betacellulin & epiregulin. Cellular Localization: Cell surface.
Formulation:	10mM PBS State: Purified State: Liquid purified IgG fraction from Bioreactor Concentrate Stabilizer: 0.05% BSA Preservative: 0.05% Sodium Azide
Concentration:	lot specific
Purification:	Protein A/G Chromatography
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~170 kDa (wild type) and ~145 kDa (vIII variant)



[View online »](#)

Gene Name:	epidermal growth factor receptor
Database Link:	Entrez Gene 1956 Human P00533
Background:	EGFR is type I receptor tyrosine kinase with sequence homology to erbB-1, -2, -3 -4 or HER-1, -2, -3 -4. It binds to Epidermal Growth Factor (EGF), Transforming Growth Factor- α (TGF- α), Heparin-binding EGF (HB-EGF), amphiregulin, betacellulin and epiregulin. EGFR is overexpressed in tumors of breast, brain, bladder, lung, gastric, head & neck, esophagus, cervix, vulva, ovary, and endometrium. It is predominantly present in squamous cell carcinomas.
Synonyms:	Epidermal growth factor receptor, EGF Receptor, erbB-1, c-ErbB-1