

Product datasheet for **AM50299PU-T**

Moesin (MSN) Mouse Monoclonal Antibody [Clone ID: MSN491]

Product data:

Product Type:	Primary Antibodies
Clone Name:	MSN491
Applications:	FC, IF, IHC, IP, WB
Recommended Dilution:	<p>ELISA: Use BSA free Antibody for coating.</p> <p>Flow Cytometry: 0.5-1 µg/million cells.</p> <p>Immunofluorescence: 0.5-1 µg/ml.</p> <p>Western Blotting: 0.5-1 µg/ml.</p> <p>Immunoprecipitation: 0.5-1 µg/500 µg protein lysate.</p> <p>Immunohistochemistry on Frozen and Formalin-Fixed Paraffin Sections: 0.5-1.0 µg/ml for 30 minutes at RT.</p> <p>Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate Buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes.</p> <p>Positive Control: HT-29, CH3LC, or HUVEC cells. Uterus, placenta, tonsil (both B and T lymphocytes), skeletal muscle, thyroid, or kidney.</p>
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Recombinant human Moesin protein.
Specificity:	<p>This Monoclonal Antibody Recognizes 78kDa Moesin protein.</p> <p>Cellular Localization: Cell surface.</p>
Formulation:	<p>10mM PBS</p> <p>State: Purified</p> <p>State: Liquid purified IgG fraction from Bioreactor Concentrate</p> <p>Stabilizer: 0.05% BSA</p> <p>Preservative: 0.05% Sodium Azide</p>
Concentration:	lot specific
Purification:	Protein A/G Chromatography
Conjugation:	Unconjugated




[View online »](#)

Storage:	Store undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	78 kDa
Gene Name:	moesin
Database Link:	Entrez Gene 4478 Human P26038


Background: Moesin, a member of the talin-4.1 superfamily, is a linking protein of the submembraneous actin cytoskeleton. It is expressed in variable amounts in cells of different phenotypes such as macrophages, lymphocytes, fibroblastic, endothelial, epithelial, and neuronal cell lines but not in blood cells. The ERM proteins, ezrin, radixin, and moesin are involved in a variety of cellular functions, such as cell adhesion, migration, and the organization of cell surface structures, and are highly homologous, both in protein sequence and in functional activity, with merlin/schwannomin, a neurofibromatosis-2-associated tumor-suppressor protein. Cell lines of epithelial and mesothelial origin contain both moesin and radixin whereas cells of endothelial and lymphoid origin express moesin.

Synonyms: MSN

Product images:



Western blot analysis of Moesin in human HT29 Cells using Moesin Antibody (Clone MSN491).



Formalin-Fixed, Paraffin-Embedded Human placenta stained with Moesin Antibody (Clone MSN491).