

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for AM50301PU-S

### EMA (MUC1) Mouse Monoclonal Antibody [Clone ID: GP1.4 + E29]

#### **Product data:**

| Product Type:         | Primary Antibodies   |
|-----------------------|--|
| Clone Name:           | GP1.4 + E29  |
| Applications:         | FC, IF, IHC, IP, WB  |
| Recommended Dilution: | <ul> <li>Flow Cytometry: 0.5-1 μg/million cells.</li> <li>Immunofluorescence: 0.5-1 μg/ml.</li> <li>Western Blotting: 0.5-1 μg/ml.</li> <li>Immunoprecipitation: 0.5-1 μg/500 μg protein lysate.</li> <li>Immunohistochemistry on Frozen and Formalin-fixed Sections: 0.5-1.0 μg/ml for 30 minutes at RT.</li> <li>No special pretreatment is required for staining of formalin/paraffin tissues.</li> <li><i>Positive Control</i>: Breast or colon carcinoma.</li> </ul>  |
| Reactivity:           | Human  |
| Host:                 | Mouse  |
| Clonality:            | Monoclonal   |
| Immunogen:            | Human milk fat globule membranes (GP1.4); Delipidated extract of human milk fat globule<br>membranes (E29).<br><b>Isotypes:</b> Mouse / IgG1, kappa (GP1.4) + Mouse / IgG2a, kappa (E29)   |
| Specificity:          | In Western blotting, it recognizes proteins in MW range of 265-400kDa, identified as different glycoforms of EMA. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhesion protein. EMA may provide a protective layer on epithelial cells against bacterial and enzyme attack. The beta subunit contains a C-terminal domain, which is involved in cell signaling, through phosphorylations and protein-protein interactions. In immunohistochemical assays, it superbly stains routine formalin/paraffin carcinoma tissues. Antibody to EMA is useful as a pan-epithelial marker for detecting early metastatic loci of carcinoma in bone marrow or liver. <i>Cellular Localization</i> : Cytoplasmic & cell surface. |
| Formulation:          | 10mM PBS<br>State: Purified<br>State: Liquid purified IgG fraction from Bioreactor Concentrate<br>Stabilizer: 0.05% BSA<br>Preservative: 0.05% Sodium Azide  |




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

# **MORIGENE** EMA (MUC1) Mouse Monoclonal Antibody [Clone ID: GP1.4 + E29] – AM50301PU-S

| Concentration:          | lot specific  |
|-------------------------|---|
| Purification:           | Protein A/G Chromatography                            |
| Conjugation:            | Unconjugated  |
| Storage:                | Store undiluted at 2-8°C.                             |
| Stability:              | Shelf life: one year from despatch.                   |
| Predicted Protein Size: | 265-400 kDa   |
| Gene Name:              | mucin 1, cell surface associated                      |
| Database Link:          | <u>Entrez Gene 4582 Human</u><br><u>P15941</u>        |
| Synonyms:               | MUC-1, PEMT, Episialin, EMA, H23AG, PUM, DF3, CA 15-3 |

## **Product images:**



Formalin-Paraffin breast carcinoma stained with CD227 / Mucin-1 / MUC1 Antibody (Clone GP1.4+E29).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US