

## Product datasheet for AM50301PU-T

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## EMA (MUC1) Mouse Monoclonal Antibody [Clone ID: GP1.4 + E29]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: GP1.4 + E29

**Applications:** FC, IF, IHC, IP, WB

**Recommended Dilution:** Flow Cytometry: 0.5-1 µg/million cells.

**Immunofluorescence:** 0.5-1 μg/ml. **Western Blotting:** 0.5-1 μg/ml.

**Immunoprecipitation:** 0.5-1 μg/500 μg protein lysate.

Immunohistochemistry on Frozen and Formalin-fixed Sections: 0.5-1.0  $\mu g/ml$  for 30

minutes at RT.

No special pretreatment is required for staining of formalin/paraffin tissues.

**Positive Control**: Breast or colon carcinoma.

Reactivity: Human Host: Mouse

Clonality: Monoclonal

Immunogen: Human milk fat globule membranes (GP1.4); Delipidated extract of human milk fat globule

membranes (E29).

**Isotypes:** Mouse / IgG1, kappa (GP1.4) + Mouse / IgG2a, kappa (E29)

**Specificity:** In Western blotting, it recognizes proteins in MW range of 265-400kDa, identified as different

glycoforms of EMA. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhesion protein. EMA may provide a protective layer on epithelial cells against bacterial and enzyme attack. The beta subunit contains a C-terminal domain, which is involved in cell signaling, through phosphorylations and protein-protein interactions. In immunohistochemical assays, it superbly stains routine formalin/paraffin carcinoma tissues. Antibody to EMA is useful as a pan-epithelial marker for detecting early metastatic loci of

carcinoma in bone marrow or liver.

*Cellular Localization*: Cytoplasmic & cell surface.

Formulation: 10mM PBS

State: Purified

State: Liquid purified IgG fraction from Bioreactor Concentrate

Stabilizer: 0.05% BSA

Preservative: 0.05% Sodium Azide





**Concentration:** lot specific

**Purification:** Protein A/G Chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 265-400 kDa


**Gene Name:** mucin 1, cell surface associated

Database Link: Entrez Gene 4582 Human

P15941

Synonyms: MUC-1, PEMT, Episialin, EMA, H23AG, PUM, DF3, CA 15-3

# **Product images:**



Formalin-Paraffin breast carcinoma stained with CD227 / Mucin-1 / MUC1 Antibody (Clone GP1.4+E29).