

Product datasheet for **AM50302PU-T**

Myogenin (MYOG) Mouse Monoclonal Antibody [Clone ID: MGN185 + F5D]

Product data:

Product Type:	Primary Antibodies
Clone Name:	MGN185 + F5D
Applications:	FC, IF, IHC, IP, WB
Recommended Dilution:	Flow Cytometry: 0.5-1 $\mu\text{g}/10^6$ cells. Immunofluorescence: 0.5-1 $\mu\text{g}/\text{ml}$. Western Blot: 0.5-1 $\mu\text{g}/\text{ml}$. Immunoprecipitation: 0.5-1 $\mu\text{g}/500$ μg protein lysate. Immunohistochemistry on Frozen and Formalin-Fixed Paraffin Sections: 0.5-1 $\mu\text{g}/\text{ml}$ for 30 minutes at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes. Positive Control: Rh-30 cells. Skeletal muscle or rhabdomyosarcoma.
Reactivity:	Feline, Human, Mouse, Porcine, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human myogenin recombinant protein (MGN185); Rat Myogenin recombinant fragment containing amino acid 30-224 (F5D). Isotypes: Mouse / IgG1, kappa (MGN185); Mouse / IgG1, kappa (F5D)
Specificity:	Anti-myogenin labels the nuclei of myoblasts in developing muscle tissue, and is expressed in tumor cell nuclei of rhabdomyosarcoma and some leiomyosarcomas. Positive nuclear staining may occur in Wilms' tumor. Cellular Localization: Nuclear.
Formulation:	10mM PBS State: Purified State: Liquid purified IgG fraction from Bioreactor Concentrate Stabilizer: 0.05% BSA Preservative: 0.05% Sodium Azide
Concentration:	lot specific
Purification:	Protein A/G Chromatography




[View online »](#)

Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	34 kDa
Gene Name:	myogenin (myogenic factor 4)
Database Link:	Entrez Gene 4656 Human P15173

Background: Myogenin is a member of the MyoD family of myogenic basic helix-loop-helix (bHLH) transcription factors that also includes MyoD, Myf-5, and MRF4 (also known as herculinor Myf-6). Family members are expressed exclusively in skeletal muscle and play a key role in activating myogenesis by binding to enhancer sequences of muscle-specific genes. The regulatory domain of MyoD is approximately 70 amino acids in length and includes both a basic DNA binding motif and a bHLH dimerization motif. Family members share about 80% amino acid homology in their bHLH motifs. Transfection of myogenin and other family members into a variety of non-muscle cells has been shown to either convert these cells to myogenic cells, or to transcriptionally activate a set of otherwise unexpressed muscle-specific genes. In addition to activating muscle specific genes, family members activate their own transcription and transactivate the transcription of other family members. For example, transfection of myogenin into 10T1/2 cells or Swiss 3T3 cells results in the activation of the endogenous myogenin gene as well as transactivation of MyoD. Likewise, the transfection of MyoD into these cells results in the activation of MyoD as well as the transactivation of myogenin. Each member of the family has distinct roles in muscle development; myogenin plays a key role in muscle maturation. Myogenin migrates at a molecular weight of ~34 kDa by SDS-PAGE.

Synonyms: MYOG, MYF4, Myogenic factor 4, BHLHC3

Product images:



Formalin-Fixed, Paraffin-Embedded Human rhabdomyosarcoma stained with Myogenin Antibody (Clone MGN185+F5D).