

## Product datasheet for **AM50317PU-T**

### TNF alpha (TNF) Mouse Monoclonal Antibody [Clone ID: TNF706]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	TNF706
Applications:	FC, IF, IHC
Recommended Dilution:	<b>ELISA:</b> Use BSA free Antibody for coating. <b>Flow Cytometry:</b> 0.5-1 µg/million cells. <b>Immunofluorescence:</b> 1-2 µg/ml. <b>Immunohistochemistry on Formalin-Fixed Paraffin Sections:</b> 1-2 µg/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes. <b>Positive Control:</b> HeLa, HL-60, or A431 cells. Macrophages in lymph node or Tonsil.
Reactivity:	Canine, Feline, Human, Mouse, Rabbit, Rat, Zebrafish
Host:	Mouse
Isotype:	IgM
Clonality:	Monoclonal
Immunogen:	Recombinant N-terminal fragment of Human TNF-alpha.
Specificity:	Recognizes TNF-alpha. <b>Cellular Localization:</b> Cytoplasmic and extracellular (secreted).
Formulation:	10mM PBS State: Purified State: Liquid purified IgG fraction from Bioreactor Concentrate Stabilizer: 0.05% BSA Preservative: 0.05% Sodium Azide
Concentration:	lot specific
Purification:	Protein A/G Chromatography
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.



[View online »](#)

**Predicted Protein Size:** 17 kDa

**Gene Name:** tumor necrosis factor

**Database Link:** [Entrez Gene 7124 Human P01375](#)

**Background:** Tumor Necrosis Factor Alpha (TNF alpha) is a protein secreted by lipopolysaccharide-stimulated macrophages, and causes tumor necrosis when injected into tumor bearing mice. TNF alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF alpha exists as a multimer of two, three, or five non-covalently linked units, but shows a single 17kDa band following SDS PAGE under non-reducing conditions. TNF alpha is closely related to the 25kDa protein Tumor Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF alpha causes cytolysis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. TNF alpha is currently being evaluated in treatment of certain cancers and AIDS Related Complex.

**Synonyms:** TNF, TNF-a, TNFA, TNFSF2, Cachectin