

Product datasheet for AM50346PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NCR2 Mouse Monoclonal Antibody [Clone ID: AT1G6]

Product data:

Product Type: Primary Antibodies

Clone Name: AT1G6

Applications: ELISA, FC, IF, WB

Recommended Dilution: The antibody has been tested by ELISA, Western blot analysis, Flow cytometry and ICC/IF to

assure specificity and reactivity. Since application varies, however, each investigation should

be titrated by the reagent to obtain optimal results.

Recommended starting dilution is 1:1000.

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Recombinant human NKp44 (19-130aa) purified from E. coli.

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein-A affinity chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: natural cytotoxicity triggering receptor 2

Database Link: Entrez Gene 9436 Human

<u>095944</u>

Background: NKp44 (NCR2) is a member of the natural cytotoxicity receptor (NCR) family that is expressed

on activated human NK cells. NKp44 displays a single extracellular Ig-like V domain and a transmembrane portion containing the charged residue (Lysine), likely involved in the

association with KARAP/DAP12 molecules.






Synonyms: NCR2, LY95, NKp44

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Natural killer cell mediated cytotoxicity

Product images:



The Human recombinant proteins, NKp30, NKp44, and NKp46 (each 20ng per well) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human NKp44 antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.