

Product datasheet for AM50630PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GNMT Mouse Monoclonal Antibody [Clone ID: AT5D1]

Product data:

Product Type: Primary Antibodies

Clone Name: AT5D1

Applications: ELISA, IF, WB

Recommended Dilution: The antibody has been tested by ELISA, Western blot analysis and ICC/IF to assure specificity

and reactivity.

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Recombinant human GNMT (1-295aa) purified from E. coli.

Formulation: Phosphate-Buffered Saline (pH 7.4) with 0.02% Sodium Azide, 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Affinity Chromatography on Protein A

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: glycine N-methyltransferase

Database Link: Entrez Gene 27232 Human

Q14749

Background: Ras-related protein Glycine N-methyltransferase (GNMT) catalyzes the methylation of glycine

by using S-adenosylmethionine (AdoMet) to form N-methylglycine (sarcosine) with the concomitant production of S-adenosylhomocysteine (AdoHcy). Possible crucial role in the

regulation of tissue concentration of AdoMet and of metabolism of methionine.

Synonyms: Glycine N-methyltransferase







Protein Families: Druggable Genome

Protein Pathways: Glycine, serine and threonine metabolism

Product images:





The Recombinant protein (50ng) and Tissue lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human GNMT antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1. GNMT recombinant protein. Lane 2: Liver tissue lysate. The Cell lysates (10ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human GNMT antibody (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. Lane 1: 293T cell lysate, Lane 2: GNMT Transfected 293T cell lysate.

ICC/IF analysis of GNMT in HepG2 cells line, stained with DAPI (Blue) for nucleus staining and monoclonal anti-human GNMT antibody (1:100) with goat anti-mouse IgG-Alexa fluor 488 conjugate (Green).