

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AM60041PU-S

SIGLEC5 / SIGLEC14 Mouse Monoclonal Antibody [Clone ID: 1A5]

Product data:

Product Type:	Primary Antibodies
Clone Name:	1A5
Applications:	ELISA, FC, IF, IP
Recommended Dilution:	Flow cytometry: 1/25-1/200, use 10µl of this working dilution to label 10e6 cells in 100µl. Immunofluorescence. Immunoprecipitation. ELISA.
Reactivity:	Chimpanzee, Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	SIGLEC5-Fc protein, consisting of the full-length extracellular region of human SIGLEC5, fused with the Fc region of human IgG1
Specificity:	This antibody recognises human SIGLEC5 / CD170. Clone 1A5 antibody is one of several SIGLEC5 antibodies which also recognises human SIGLEC14 (<i>Angata, T. et al., 2006</i>).
Formulation:	PBS containing 0.09% Sodium azide State: Purified State: Liquid purified IgG fraction
Concentration:	lot specific
Purification:	Affinity chromatography on Protein G
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

SIGLEC5 / SIGLEC14 Mouse Monoclonal Antibody [Clone ID: 1A5] – AM60041PU-S

Background:Siglec-5 shares an almost identical sequence with Siglec-14 within the first two Ig-like
domains, indicating partial gene conversion between these two Siglecs, also evident in other
primate species.

Siglec-5, also known as CD170, is a sialic-acid-binding Ig-like lectin, and member of the Ig superfamily, expressed by dendritic cells (DCs), activated macrophages, neutrophils, and cells of the monocyte/myeloid lineage. Highly related to the myelomonocytic-derived adhesion molecule CD33 (Siglec-3), Siglec-5 mediates sialic-acid dependent binding to cells, and is as well acting as an inhibitory receptor in the down-regulation of cell activation. Structurally, Siglec-5 contains an immunoreceptor tyrosine-based inhibitor motif (ITIM), which plays a part in the modulation of cellular responses, and when phosphorylated, can bind to the SH2 domain of several SH2-containing phosphatases.

Siglec-14 is a putative sialic-acid binding adhesion molecule and predominantly expressed in hematopoietic tissues, which have been shown to associate with the activating adapter protein DAP12.

Synonyms: Q08ET2, O15389, CD170

Product images:



Human peripheral blood monocytes stained with SIGLEC5 / SIGLEC14 antibody Cat.-No. AM60041PU-S followed by secondary FITC conjugated antibody.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US