

Product datasheet for AM60044FC-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SIGLEC10 Mouse Monoclonal Antibody [Clone ID: 5G6]

Product data:

Product Type: Primary Antibodies

Clone Name: 5G6
Applications: FC

Recommended Dilution: Flow cytometry: Neat - 1/10, use 10ul of the suggested working dilution to label 10e6 cells.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Recombinant human SIGLEC10, fused with the Fc region of human IgG

Specificity: This antibody clone 5G6 specifically recognizes human SIGLEC10.

Studies have shown that clone 5G6 does not cross react with SIGLECs 3, 5, 7, 8 and 9Munday

J. et al.)

Formulation: PBS containing 0.09% Sodium azide and 1% BSA

Label: FITC State: Purified

State: Liquid purified IgG

Label: Fluorescein Isothiocyanate Isomer 1

Concentration: lot specific

Purification: Affinity chromatography on Protein G

Conjugation: FITC

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: sialic acid binding Ig like lectin 10

Database Link: Entrez Gene 89790 Human

Q96LC7






Background:

Siglec10 (Sialic acid binding Ig like lectin 10), a putative adhesion molecule and member of the Ig superfamily, is expressed by monocytes, B cells, eosinophils, and at a higher level by a subpopulation of CD16+CD56- natural killer (NK) cells.

Structurally, Siglec10 is most similar to the CD33 related group of Siglecs, and preferentially binds to glycoconjugates containing alpha 2,3 or alpha2,6 linked sialic acid. Studies have shown that Siglec10 acts as a substrate for VAP1 (Vascular adhesion protein1), a glycoprotein expressed on endothelium during inflammation, which is involved in primary amine oxidation and leucocyte trafficking (*Kive, E. et al.*). This interaction between Siglec10 and VAP1, implicates Siglec10 in endothelial lymphocyte adhesion and in the modulation of the inflammatory microenvironment.

Synonyms: Siglec-10, SLG2

Product images:



Human peripheral blood lymphocytes stained with mouse anti-human siglec-10-FITC clone 5G6.