

## Product datasheet for AP00766PU-N

## **Fibrinogen Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, WB

Recommended Dilution: ELISA.

Western Blot.

Reactivity: Mouse Host: Rabbit

Clonality: Polyclonal

Immunogen: Native Mouse Fibrinogen.

**Specificity:** This antibody recognizes Mouse Fibrinogen.

Cross reacts with Fibrin.

Formulation: 0.05M Sodium Phosphate, pH 6.6 containing 0.1M Sodium Chloride and 1 mM EDTA

State: Purified

State: Liquid purified Ig fraction

Preservative: None

**Concentration:** lot specific

**Purification:** Protein A Chromatography

Conjugation: Unconjugated

Storage: Upon receipt, store (in aliquots) at -20°C to -80°C.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## Fibrinogen Rabbit Polyclonal Antibody - AP00766PU-N

Background:

Fibrinogen is the main protein of blood coagulation system. It is a large protein and it consists of two identical subunits that contain three polypeptide chains: alpha, beta and gamma. All chains are connected with each other by a number of disulfide bonds. Fibrinopeptides A (1 to 16 amino acids) and B (1 to 17 amino acids) are released by thrombin from the N terminal parts of alpha and beta chains, respectively. In this way fibrinogen is converted into fibrin, which by means of polymerization forms a fibrin clot. Fibrinogen clotting underlies pathogenesis of MI, thromboembolism and thromboses of arteries and veins, since fibrin is the main substrate for thrombus formation. Fibrinogen activation is also involved in pathogenesis of inflammation, tumor growth and many other diseases. The normal fibrinogen concentration in plasma is about 3 mg/ml. The elevated level of fibrinogen in patient's blood is regarded as an independent risk factor for cardiovascular diseases. An increase in blood fibrinogen concentration was shown to be a strong predictor of coronary heart disease (Sonel A. et al, and Rapold H.J. et al). All these facts make fibrinogen an important parameter in the diagnosis of cardiovascular diseases.

**Synonyms:** FGA, FGB, FGG