

## Product datasheet for **AP01114BT-N**

### **RANKL (TNFSF11) Goat Polyclonal Antibody**

#### **Product data:**

<b>Product Type:</b>	Primary Antibodies
<b>Applications:</b>	ELISA, WB
<b>Recommended Dilution:</b>	ELISA: Direct: To detect hsRANKL by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml is required. In conjunction with compatible secondary reagents, it allows the detection of at least 0.2 - 0.4 ng/well of recombinant hsRANKL. Sandwich: To detect hsRANKL by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25 - 1.0 µg/ml is required. In conjunction with Polyclonal Anti-Human sRANKL as a capture antibody, it allows the detection of at least 0.2 - 0.4 ng/well of recombinant hsRANKL. Western Blot: To detect hsRANKL by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hsRANKL is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
<b>Reactivity:</b>	Human
<b>Host:</b>	Goat
<b>Clonality:</b>	Polyclonal
<b>Immunogen:</b>	Highly pure (> 98 %) recombinant human sRANKL
<b>Specificity:</b>	This antibody detects RANKL.
<b>Formulation:</b>	PBS, pH 7.2 Label: Biotin State: Sterile filtered lyophilized Ig fraction
<b>Reconstitution Method:</b>	Centrifuge vial prior to opening. Restore in sterile PBS containing 0.1 % BSA to a concentration of 0.1 - 1.0 mg/ml.
<b>Purification:</b>	Affinity chromatography
<b>Conjugation:</b>	Biotin
<b>Storage:</b>	Store the lyophilized antibody at -20 °C. Following reconstitution it is stable for two weeks at 2 - 8 °C. Frozen aliquots are stable for 6 months when stored at -20 °C. Avoid repeated freezing and thawing.
<b>Stability:</b>	Shelf life: One year from despatch.



[View online »](#)

**Gene Name:** tumor necrosis factor superfamily member 11

**Database Link:** [Entrez Gene 8600 Human O14788](#)

**Background:** RANKL is a member of the tumor necrosis factor (TNF) cytokine family which is a ligand for osteoprotegerin and functions as a key factor for osteoclast differentiation and activation. There are three isoforms of RANKL. Human RANKL is a soluble 20 kDa polypeptide, comprising the TNF homologous region of RANKL (176 amino acid residues). This protein was shown to be a dendritic cell survival factor and is involved in the regulation of T cell dependent immune response. T cell activation was reported to induce expression of this gene and lead to an increase of osteoclastogenesis and bone loss. This protein was shown to activate antiapoptotic kinase AKT/PKB through a signaling complex involving SRC kinase and tumor necrosis factor receptor associated factor (TRAF) 6, which indicated that this protein may have a role in the regulation of cell apoptosis. RANKL deficient mice show severe osteoporosis and complete absence of osteoclasts as a result of lack of osteogenesis.

**Synonyms:** OPGL, RANK Ligand, RANKL, TRANCE, TNFSF11, ODF