

Product datasheet for **AP01138PU-S**

IGFBP3 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	Sandwich ELISA: To detect hIGF-BP3 (using 100 µl/well antibody solution) a concentration of 0.5-2.0 µg/ml of this antibody is required. In conjunction with Biotinylated Anti-Human IGF-BP3 as a detection antibody, it allows the detection of at least 0.2-0.4 ng/well of recombinant hIGF-BP3. Western blot: To detect hIGF-BP3 this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIGF-BP3 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions. Immunohistochemistry on Formalin-Fixed, Paraffin Sections: 0.5-2.5 µg/ml for 2 hours at RT. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 Sodium Citrate buffer is recommended. <i>Positive Control:</i> Normal Human Placenta.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure (> 98%) recombinant Human IGF-BP3
Specificity:	This antibody detects Insulin-like Growth Factor Binding Protein 3.
Formulation:	PBS, pH 7.2 State: Aff - Purified State: Lyophilized (sterile filtered) Ig fraction
Reconstitution Method:	Centrifuge vial prior to opening. Restore in sterile water to a concentration of 0.1-1.0 mg/ml.
Purification:	Immunoaffinity chromatography
Conjugation:	Unconjugated
Storage:	Store the lyophilized antibody at -20°C. Following reconstitution it is stable for two weeks at 2-8°C. Frozen aliquots are stable for 6 months when stored at -20 °C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.



[View online »](#)

Gene Name: insulin like growth factor binding protein 3

Database Link: [Entrez Gene 3486 Human P17936](#)

Background: Insulin like growth factor binding protein 3 (IGFBP3) is a member of the superfamily of insulin like growth factor (IGF) binding proteins which include six high affinity IGF binding proteins (IGFBP) and at least four low affinity binding proteins referred to as IGFBP related proteins (IGFBPrP). The IGFBP members are cysteine rich proteins with conserved cysteine residues clustered in the amino terminal and the carboxy terminal regions of the molecule. The cDNA sequence encoding the mature human IGFBP3 is fused to the signal peptide of CD33. Human IGFBP3 is the major IGF binding protein in plasma where it exists in a ternary complex with IGFI or IGFI and an acid labile subunit. IGFBPs hold a central position in IGF ligand-receptor interactions through influences on both the bioavailability and distribution of IGFs in the extracellular environment. Insulin like growth factor binding protein 3 (IGFBP3) can modulate the mitogenic and metabolic effects of the insulin like growth factors (IGFs). Insulin like growth factor binding protein 3 (IGFBP3) is expressed in multiple tissues. The highest expression level is found in the non paranchymal cells of the liver. The expression levels are higher during extrauterine life and peak during puberty. Insulin like growth factors (IGFs) and IGF binding proteins (IGFBPs) play important roles in cell growth and differentiation. IGFBP3 is one of the factors in serum that is responsible for high serum induced apoptosis in PC3 cells, a prostate cancer cell line. IGFBP3 is important in controlling glucose homeostasis with increased urinary levels in type I diabetes with persistent microalbuminuria.

Synonyms: IGFBP-3, IGF-BP3, IBP3, Insulin-like growth factor-binding protein 3, IGF-binding protein 3, IBP-3