

## Product datasheet for **AP01153PU-N**

### MIA2 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	<b>Indirect ELISA:</b> To detect hMIA-2 (using 100µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. In conjunction with compatible secondary reagents, it allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIA-2. <b>Sandwich ELISA:</b> To detect hMIA-2 (using 100µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. In conjunction with Biotinylated Anti-Human MIA-2 as a detection antibody, it allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIA-2. <b>Western blot:</b> To detect hMIA-2 this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMIA-2 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Highly pure (> 98 %) E.coli derived 11.5 kDa recombinant human MIA-2
Specificity:	This antibody detects MIA-2.
Formulation:	PBS, pH 7.2 without preservatives. State: Aff - Purified State: Sterile filtered lyophilized Ig fraction
Reconstitution Method:	Centrifuge vial prior to opening. Restore in sterile water to a concentration of 0.1 - 1.0 mg/ml.
Purification:	Immunoaffinity chromatography
Conjugation:	Unconjugated
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.



[View online »](#)

**Gene Name:** CTAGE family member 5

**Database Link:** [Entrez Gene 4253 Human Q96PC5](#)

**Background:** MIA2 (Melanoma inhibitory activity 2) is a secreted cytokine and a member of the MIA/OTOR family. It is predominantly expressed in hepatocytes and may play a role in the pathophysiology of liver disease. Elevated levels of MIA2 may represent a clinically useful marker for diagnosis of hepatic disease activity and severity. MIA2 may act as a tumor suppressor in hepatocellular carcinoma.

**Synonyms:** Melanoma inhibitory activity protein 2