

Product datasheet for **AP01170PU-S**

RPL10 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	Western Blot: 1/500 - 1/1000.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to the N-terminal of Human RPL10.
Specificity:	This antibody detects endogenous levels of QM protein. (region surrounding Phe34)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch
Predicted Protein Size:	~ 25 kDa
Gene Name:	ribosomal protein L10
Database Link:	Entrez Gene 6134 Human P27635




[View online »](#)

Background:

The c-Jun protein is a major component of the transcription factor AP-1, originally shown to mediate phorbol ester tumor promoter (TPA)-induced expression of responsive genes through the TPA-response element (TRE). The Jun proteins form homo- and heterodimers which bind the TRE, while Fos proteins are active only as heterodimers with any of the Jun proteins. Fos/Jun heterodimers have a much higher affinity for the TRE than Jun homodimers. A distant member of the MAP kinase family, designated c-Jun NH2-terminal kinase (JNK1) functions to regulate c-Jun by phosphorylation at the amino terminal serine regulatory sites, Ser 63 and Ser 73). QM has been described as a transcription factor that can function to bind DNA directly or alternatively can interact with c-Jun to inhibit transactivation of AP-1 promoter driven reporter vectors by Jun-Jun homodimers. QM is highly conserved throughout eukaryotic evolution and is apparently a member of a multi-gene family.

Synonyms:

DXS648E, QM

Product images:

Western blot (WB) analysis of QM antibody in extracts from K562 cells treated with Insulin 0.01 u/ml 15 min.