

## Product datasheet for AP01183PU-M

# ST14 Rabbit Polyclonal Antibody

**Product data:** 

**Product Type:** Primary Antibodies

Applications: IF, WB

Recommended Dilution: Western Blot: 1/500 - 1/1000.

Immunofluorescence: 1/50 - 1/200.

Reactivity: Human

Host: Rabbit

Clonality: Polyclor

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide, corresponding to the N-terminual of Human Matriptase.

**Specificity:** This antibody detects endogenous levels of Matriptase protein. (region surrounding Lys21)

**Formulation:** Phosphate buffered saline (PBS), pH~7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE).

Preservative: 15 mM Sodium Azide

Concentration: 1.0 mg/ml

**Purification:** Affinity Chromatography using epitope-specific immunogen.

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: One year from despatch

Predicted Protein Size: ~ 90 kDa

**Gene Name:** suppression of tumorigenicity 14

Database Link: Entrez Gene 6768 Human

Q9Y5Y6



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




### Background:

Matriptase (also known as MT-SP1, ST14, prostamin and epithin in mouse) is a tumorassociated type II transmembrane serine protease that is highly expressed in many human cancer-derived cell lines and is implicated in extracellular matrix re-modeling, tumor growth, and metastasis. Matriptase performs pleiotropic functions in the development of the epidermis, hair follicles, and cellular immune system. Sphingosine 1-phosphate (S1P, SPP), present in serum-derived lipoproteins, activates matriptase while matriptase activates both urokinase-type plasminogen activator and hepatocyte growth factor (HGF). Hepatocyte growth factor activator inhibitor type 1 (HAI-1) is a Kunitz-type serine protease inhibitor identified as a strong inhibitor of matriptase and HGF. Advanced-stage ovarian tumors that express matriptase are more likely to do so in the absence of its inhibitor, HAI-1, indicating that an imbalance in the matriptase: HAI-1 ratio could be important in the development of advanced disease.

Synonyms:


PRSS14, SNC19, TADG15, T-SP1, Prostamin, Serine protease 14, Serine protease TADG-15

# **Product images:**



Western blot (WB) analysis of Matriptase antibody in extracts from A549.





Hela whole cell lysate Matriptase (K21) pAb at 1:500 dilution Western blot (WB) analysis of Matriptase antibody in extracts from hela cells.