

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AP01190PU-M

Adenosine A3 Receptor (ADORA3) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies	
Applications:	ELISA, IF, IHC, WB	
Recommended Dilution:	Western Blot: 1/500-1/1000. Immunofluorescence: 1/50-1/200. Immunohistochemistry on Paraffin Sections: 1/50-1/200.	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Immunogen:	Synthetic peptide, corresponding to amino acids C-terminus of Human ADORA3.	
Specificity:	This antibody detects endogenous levels of Adenosine A3-R protein. (region surrounding Glu318)	
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE) Preservative: 0.05% Sodium Azide	
Concentration:	1.0 mg/ml	
Purification:	Affinity Chromatography using epitope-specific immunogen	
Conjugation:	Unconjugated	
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.	
Stability:	Shelf life: One year from despatch.	
Predicted Protein Size:	~38 kDa	
Gene Name:	adenosine A3 receptor	
Database Link:	<u>Entrez Gene 140 Human</u> <u>P0DMS8</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Adenosine A3 Receptor (ADORA3) Rabbit Polyclonal Antibody – AP01190PU-M
Background:	Adenosine is involved in a variety of processes, including the synthesis of urea, the anti- inflammatory response and the inhibition of protein synthesis The adenosine receptors, including adenosine A1-R, adenosine A2A-R, adenosine A2B-R and adenosine A3-R, are integral membrane proteins that are members of the G protein-coupled receptor family. The A1-R protein mediates ureagenesis in a partially calcium-dependent manner. Adenosine is known to mediate coronary vasodilation via the A2A-R receptor. Collagen synthesis and total protein synthesis are inhibited in certain cells by adenosine, acting via the A2B receptors. Activation of the A3-R receptor inhibits the induction of the cytokine TNFalpha and blocks the endotoxin CD14 receptor signal transduction pathway.

Synonyms:	ADORA3, Adenosine A3 receptor
Synonyms.	ADDIAS, Additional AS receptor

Product images:



Western blot analysis of Adenosine A3-R Antibody in extracts from HepG2 cells.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US