

## **Product datasheet for AP01259PU-N**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Adenosine Receptor A2a (ADORA2A) Rabbit Polyclonal Antibody

**Product data:** 

Product Type: Primary Antibodies

Applications: ELISA, IF, IHC, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunofluorescence: 1/50-1/200.

**Immunohistochemistry on Paraffin Sections:** 1/50-1/200.

Reactivity: Human

Host: Rabbit

Clonality: Polyclonal

**Immunogen:** Synthetic peptide, corresponding to amino acids 330-380 of Human ADORA2A.

**Specificity:** This antibody detects endogenous levels of Adenosine A2A-R protein. (region surrounding

Tyr361)

**Formulation:** Phosphate buffered saline (PBS), pH~7.2

State: Aff - Purified

State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE)

Preservative: 0.05% Sodium Azide

Concentration: 1.0 mg/ml

**Purification:** Affinity Chromatography using epitope-specific immunogen

**Conjugation:** Unconjugated

**Storage:** Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** ~45 kDa

Gene Name: adenosine A2a receptor

Database Link: Entrez Gene 135 Human

P29274






#### Background:


Adenosine is involved in a variety of processes, including the synthesis of urea, the anti-inflammatory response and the inhibition of protein synthesis. The adenosine receptors, including Adenosine A1-R, Adenosine A2A-R, Adenosine A2B-R and Adenosine A3-R, are integral membrane proteins that are members of the G protein-coupled receptor family. The A1-R protein mediates ureagenesis in a partially calcium-dependent manner. Adenosine is known to mediate coronary vasodilation via the A2A-R receptor. Collagen synthasis and total protein synthesis are inhibited in certain cells by adenosine, acting via the A2B receptors. Activation of the A3-R receptor inhibits the induction of the cytokine TNF $\alpha$  and blocks the endotoxin CD14 receptor signal transduction pathway.

Synonyms: ADORA2A, ADORA2

# **Product images:**



Western blot analysis of Adenosine A2A-R Antibody in extracts from HepG2 cells.



Immunohistochemistry analysis of Adenosine A2A-R Antibody in paraffin-embedded human skeleta I muscle tissue.