

Product datasheet for **AP01271PU-N**

HRH1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IF, WB
Recommended Dilution:	Western blot: 1/500-1/1000. Immunofluorescence: 1/50-1/200.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 150-200 of Human Histamine H1 Receptor.
Specificity:	This antibody detects endogenous levels of Histamine H1 Receptor protein. (region surrounding Lys179)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE). Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~60 kDa
Gene Name:	histamine receptor H1
Database Link:	Entrez Gene 3269 Human P35367




[View online »](#)

Background:

Histamine is an inflammatory mediator that is ubiquitously expressed and has a broad range of pharmacologic effects. Specifically, it plays a role in the central nervous, gastrointestinal, respiratory and immune systems. The effects of histamine are mediated by a family of G protein-coupled receptors, the Histamine H1, H2, H3 and H4 Receptors. The gene encoding the human Histamine H1 Receptor maps to chromosome 3p25 and is expressed in highest abundance in placenta, with lower levels in lung, skeletal muscle, kidney and brain. The murine Histamine H2 Receptor gene maps to chromosome 13 and is highly expressed in stomach with moderate expression in brain and heart. The gene encoding the human Histamine H3 Receptor is located on chromosome 20 and is expressed as six alternative splice variants in thalamus. The human Histamine H4 Receptor gene maps to chromosome 18q11 and is expressed most abundantly in bone marrow and spleen in addition to peripheral blood leukocytes, thymus, small intestine and colon.

Synonyms:

HRH1, HR-H1, Histamine H1 receptor

Product images:

Western blot (WB) analysis of Histamine H1 Receptor antibody in extracts from COLO205 cells.