

## Product datasheet for **AP01280PU-M**

### ID4 (N-term) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	<b>Western blot:</b> 1/500-1/1000. <b>Immunohistochemistry on Paraffin sections:</b> 1/50-1/200.
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids N-terminus of Human Id4.
Specificity:	This antibody detects endogenous levels of ID4 / BHLHB27 protein (region surrounding Ser5).
Formulation:	Phosphate buffered saline (PBS), pH 7.2 State: Aff - Purified State: Liquid purified Ig fraction Preservative: 0.05% Sodium azide
Concentration:	1.0 mg/ml
Purification:	Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE)
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 17 kDa
Gene Name:	inhibitor of DNA binding 4, HLH protein
Database Link:	<a href="#">Entrez Gene 3400 Human P47928</a>


[View online »](#)

**Background:**

Transcription factors containing a basic helix-loop-helix (bHLH) motif regulate the expression of tissue-specific genes in a number of mammalian and insect systems. DNA-binding activity of the bHLH proteins is dependent on formation of homo- and/or heterodimers. Dominant-negative HLH proteins (encoded by Id-related genes) also contain the HLH-dimerization domain but lack the DNA-binding basic domain. Consequently, Id proteins inhibit binding to DNA and transcriptional transactivation by heterodimerization with bHLH proteins (referenced from entrez gene).

**Synonyms:**

DNA-binding protein inhibitor ID-4, Inhibitor of DNA binding 4

**Product images:**


Western blot analysis with extracts from HepG2 cells using ID4 / BHLHB27 antibody.