

Product datasheet for **AP01287PU-N**

IRAKM (IRAK3) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	Western Blot: 1/500-1/1000. Immunohistochemistry on Paraffin Sections: 1/50-1/200. Immunofluorescence: 1/50-1/200.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 500-540 of Human IRAK-M.
Specificity:	This antibody detects endogenous levels of IRAK-M protein. (region surrounding Gln649)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% by SDS-PAGE). Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity-Chromatography using epitope-specific immunogen.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~68 kDa
Gene Name:	interleukin 1 receptor associated kinase 3
Database Link:	Entrez Gene 11213 Human Q9Y616




[View online »](#)

Background:


Interleukin-1 receptor (IL1R)-associated kinases (IRAKs) are important mediators in the signal transduction of Toll-like receptor (TLR) and IL1R family members, collectively referred to as TIRs. Binding of IL-1 to its cognate receptor results in the activation of the NFκB signaling pathway. A 100 kDa IL-1- dependent kinase termed IRAK-1 (for IL-1 receptor- associated kinase 1) coimmunoprecipitates with activated IL-1RI and is implicated as an upstream mediator of NFκB activation. A related *Drosophila* protein, Pelle, is a known upstream activator of Dorsal, the *Drosophila* homolog of NFκB. IRAK-2 is a proximal mediator of IL1, a component of the IL1R signaling complex, and is required for IL1R-induced NFκB activation. IRAK-4, like IRAK-1 and Pelle, has auto- and cross-phosphorylation kinase activity. IRAK-4 is strongly expressed in kidney and is also found in lung, testis, small intestine, breast, liver, and placenta. In contrast to the other IRAKs that are expressed in most cell types, IRAK-M is restricted to monocytic cells. IRAK-M mRNA transcripts are found predominantly in PBL and the monocytic cell lines U937 and THP-1.

Synonyms:

IRAK-3, IRAK-M

Product images:


Western blot (WB) analysis of IRAK-M antibody in extracts from K562 cells.



Immunohistochemistry (IHC) analyzes of IRAK-M antibody in paraffin-embedded human brain tissue.