

Product datasheet for AP01299PU-M

Frizzled 7 (FZD7) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IF, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunofluorescence: 1/50-1/200.

Reactivity: Human, Mouse

Host: Rabbit

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 61-110 of Human Frizzled-7.

Specificity: This antibody detects endogenous levels of Frizzled-7 protein. (region surrounding Tyr87)

Formulation: Phosphate buffered saline (PBS), pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction. Preservative: 0.05% Sodium Azide

Concentration: 1.0 mg/ml

Purification: Affinity Chromatography.

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 64, 85 kDa

Gene Name: frizzled class receptor 7

Database Link: Entrez Gene 8324 Human

O75084



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The frizzled gene, originally identified in Drosophila melanogaster, is involved in the development of tissue polarity. The mammalian homolog of frizzled, as well as several secreted mammalian frizzled-related proteins: FRP-1 (also designated SARP2), FRP-2 (also designated SARP1), FRP-3, FRP-4 and SARP3 (also designated FRP-5), have been identified. The frizzled proteins contain seven transmembrane domains, a cysteine-rich domain in the extracellular region and a carboxy terminal Ser/Thr-xxx-Val motif. They function as receptors for Wnt. frizzled-7 has a Lys-Thr-X-X-X-Trp motif which is involved in the activation of the Wnt/beta-catenin signaling pathway. frizzled-7 is an integral membrane protein showing a high expression in adult skeletal muscle, fetal kidney, fetal lung, adult heart, brain and placenta. It is specifically expressed in squamous cell esophageal carcinomas.

Synonyms: Fz-7, mFz7

Product images:



Western blot (WB) analysis of Frizzled-7 antibody in extracts from HT-29 cells.