

## Product datasheet for **AP01307PU-N**

### **BAI1 (ADGRB1) Rabbit Polyclonal Antibody**

#### **Product data:**

Product Type:	Primary Antibodies
Applications:	ELISA, IF, IHC, WB
Recommended Dilution:	<b>Western Blot:</b> 1/500-1/1000. <b>Immunofluorescence:</b> 1/50-1/200. <b>Immunohistochemistry on paraffin sections:</b> 1/50-1/200.
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 711-760 of Human BAI-1.
Specificity:	This antibody detects endogenous levels of BAI-1 protein. (region surrounding Gln732)
Formulation:	Phosphate buffered saline (PBS), pH~7.2 State: Aff - Purified State: Liquid purified Ig fraction (> 95% pure by SDS-PAGE) Preservative: 0.05% Sodium Azide
Concentration:	1.0 mg/ml
Purification:	Affinity Chromatography using epitope-specific immunogen
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	~ 174 kDa
Gene Name:	adhesion G protein-coupled receptor B1
Database Link:	<u><a href="#">Entrez Gene 575 Human</a></u> <u><a href="#">Q14514</a></u>




[View online »](#)

**Background:**


Brain-specific angiogenesis inhibitors, including BAI-1, BAI-2 and BAI-3, are integral membrane proteins belonging to the G protein-coupled receptor 2 family. In addition to inhibiting angiogenesis in the brain, BAI proteins are also expressed in the heart, thymus, skeletal muscle, and a variety of cell lines. BAI-1 protein is specifically expressed in the brain and found to localize to the cytoplasm and membrane in neuronal cells of the cerebral cortex. Reduced expression of BAI-1 in some glioblastoma cell lines and cancer tissues implicates the functional role of BAI-1 as an inhibitor of angiogenesis. The exact mechanisms underlying BAI-1 anti-angiogenetic activity are still being investigated. BAI-1 may be involved in mediating the p53 signal in suppression of glioblastoma, as well as in cell adhesion and signal transduction. Additional research shows an inverse correlation with vascularization and BAI-1 expression in both colorectal carcinomas and pulmonary adenocarcinomas.

**Synonyms:**

BAI-1

**Product images:**


Western blot (WB) analysis of BAI-1 antibody in extracts from Jurkat treated with HepG2.



Immunohistochemistry (IHC) analyzes of BAI-1 antibody in paraffin-embedded human brain.