

Product datasheet for AP01325PU-M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Neurotensin Receptor 1 (NTSR1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Western Blot: 1/500-1/1000.

Immunofluorescence: 1/50-1/200.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Reactivity: Human, Mouse

Host: Rabbit

Clonality: Polyclonal

Specificity: NTR1 antibody detects endogenous levels of NTR1 protein. (region surrounding His218)

Formulation: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

State: Aff - Purified

State: Liquid purified Ig fraction

Concentration: 1.0 mg/ml

Purification: Affinity chromatography

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: ~ 36 kDa

Gene Name: neurotensin receptor 1 (high affinity)

Database Link: Entrez Gene 4923 Human

P30989




Background:

Neurotensin (NT) initiates an intracellular response by interacting with the G protein-coupled receptors NTR1 (NTS1 receptor, high affinity NTR) and NTR2 (NTS2 receptor, levocabastine-sensitive neurotensin receptor), and the type I receptor NTR3 (NTS3 receptor, sortilin-1, Gp95). NT has a wide distribution in regions of the brain and in peripheral tissues where NT receptors can contribute to hypotension, hyperglycemia, hypothermia, antinociception and regulation of intestinal motility and secretion. HL-60 cells express NTR1, which can couple to Gq, Gi/o, or Gs. Alternative splicing of rat NTR2 can generate a 5-transmembrane domain variant isoform that is co-expressed with the full-length NTR2 throughout the brain and spinal cord. NTR3 activation in the murine microglial cell line N11 induces MIP-2, MCP-1, IL-1 β and TNF α in an ERK1/2 and Akt kinase-dependent manner.

Synonyms:

NT-R-1,NTSR1, NTRR, NTR1

Product images:



Western blot (WB) analysis of NTR1 antibody (Cat.-No.: [AP01325PU-N]) in extracts from Jurkat cells.